arXiv:1106.0741v1 [math.AC] 4 Jun 2011

COHEN-MACAULAYNESS OF REES ALGEBRAS OF DIAGONAL IDEALS

KUEI-NUAN LIN

Abstract. Given two determinantal rings over a field k. We consider the Rees algebra of the diagonal ideal, the kernel of the multiplication map. The special fiber ring of the diagonal ideal is the homogeneous coordinate ring of the join variety. When the Rees algebra and the Symmetric algebra coincide, we show that the Rees algebra is Cohen-Macaulay.

1. Introduction

Determinantal rings and varieties have been a central topic of commutative algebra and algebraic geometry. The embedded join of two subschemes X, Y of $\mathbb{P}^n_k$ is another important subject. When $X = Y$, the join construction yields the classical secant variety. Join varieties are an important topic in algebraic geometry. The embedded join of X and Y is the closure of the union of all lines passing through two distinct points of X and Y. An important question is whether the vaiety is all of $\mathbb{P}^n_k$. As the special fiber ring of the diagonal ideal is the homogeneous coordinate ring of the embedded join, it is natural to investigate the blowup along the diagonal, rather than just the special fiber in the blow up.

To study join varieties of determinantal varieties, we investigate blowups in products of determinantal varieties. It turns out that for some of the cases where the embedded join is the whole space [S-U], the Rees algebra and the symmetric algebra of the diagonal ideal coincide [L]. In this work, we show that the Rees algebras are Cohen-Macaulay in those cases. This continues work of Simis and Ulrich [S-U], and of the author [L].

We now describe the setting. Let k be a field, $2 \leq m \leq n$ integers, $X = [x_{ij}]$ an $m \times n$ matrix of variables over k, and $I = I_{u_1}(X)$, $J = I_{u_2}(X)$ the ideals of $k[X]$ generated by the $u_1 \times u_1$ minors of X and the $u_2 \times u_2$ minors of X. Let $R_1 = k[X]/I$, $R_2 = k[X]/J$ be two determinantal rings. We consider the diagonal ideal $\mathbb{D}$ of $S = R_1 \otimes_k R_2$, defined via the exact sequence

$$0 \longrightarrow \mathbb{D} \longrightarrow S \overset{\text{mult.}}{\longrightarrow} k[X]/(I + J) \longrightarrow 0.$$

The ideal $\mathbb{D}$ is generated by the images of $x_{ij} \otimes 1 - 1 \otimes x_{ij}$ in S. The homogeneous coordinate ring of the embedded join variety is the k-subalgebra of S generated by the images of $x_{ij} \otimes 1 - 1 \otimes x_{ij}$. Those elements are homogeneous of degree 1. The homogeneous coordinate ring of the embedded joint variety $\mathcal{J}(I, J) \subseteq \mathbb{P}^{mn-1}_k$ of the determinantal

---

2000 *Mathematics Subject Classification.* Primary 13C40, 14M12; Secondary 13P10, 14Q15, 05E40.

Key words and phrases. Rees Algebra, Secant Variety, Join Variety, Determinantal Ring, Symmetric Algebra, Alexander Dual, Regularity.

varieties $V(I),\ V(J)$ in $\mathbb{P}_k^{mn-1}$ can be identified with $\mathcal{R}(\mathbb{D}) \otimes_S k = \mathcal{F}(\mathbb{D})$ regarding k as $S/\mathfrak{m}$ where $\mathfrak{m}$ is the homogeneous maximal ideal of S.

The scheme $\mathrm{Proj}(\mathcal{F}(\mathbb{D}))$ is the special fiber in the blowup $\mathrm{Proj}(\mathcal{R}(\mathbb{D}))$ of $\mathrm{Spec}(S)$ along $V(\mathbb{D})$. In this work, we study, more broadly, the blowup, rather than the special fiber.

Theorem 1. *The Rees algebra $\mathcal{R}(\mathbb{D})$ is Cohen-Macaulay if I and J are generated by the maximal minors of submatrices of X.*

In [L], the defining ideals of Rees algebras of diagonal ideals have been determined in the setting of theorem 1. Let $\mathcal{K}$ be the defining ideal of the Rees algebra of $\mathbb{D}$. By the proposition below, we deduce that $\mathcal{K}$ is Cohen-Macaulay once we show $\mathrm{in}(\mathcal{K})$ is Cohen-Macaulay.

Proposition 2. (*a*) [E][15.15] *Let R be a polynomial ring over a field k, $>$ be a monomial order on R, I an ideal of R and $\mathrm{in}(I)$ the initial ideal of I with respect to the term order $>$. Let $a_1, ..., a_r$ be polynomials in R such that $\mathrm{in}(a_1), ..., \mathrm{in}(a_r)$ form a regular sequence on $R/\mathrm{in}(I)$. Then $a_1, ..., a_r$ is a regular sequence on R/I.*

(b) [E][15.16, 15.17] *If $R/\mathrm{in}(I)$ is Cohen-Macaulay, then so is R/I.*

We use combinatorial commutative algebra to show $\mathrm{in}(\mathcal{K})$ is Cohen-Macaulay. With respect to a suitable term order, $\mathrm{in}(\mathcal{K})$ is generated by square-free monomials [L]. Square-free monomial ideals in a polynomial ring are also known as Stanley–Reisner ideals. This leads us to consider Alexander dual ideals:

Theorem 3. [E-R] *Let I be a square-free monomial ideal in a polynomial ring R. The ring R/I is Cohen-Macaulay if and only if the Alexander dual ideal I^* has a linear free resolution.*

With Theorem 3, we need to show that $(\mathrm{in}(\mathcal{K}))^*$, the Alexander dual ideal of $\mathrm{in}(\mathcal{K})$, has a linear free resolution. To do so, we find a suitable filtration starting from the Alexander dual ideal of $\mathrm{in}(\mathcal{K})$.

Acknowledgments: This work is based on author's Ph. D. thesis from Purdue University under the direction of Professor Bernd Ulrich. The author is very grateful for so many useful suggestions from Professor Ulrich.

2. Defining Equations of Rees Algebras

Let k be a field, $2 \leq m \leq n$ integers, $X_{mn} = [x_{ij}]$, $Y_{mn} = [y_{ij}]$, $Z_{mn} = [z_{ij}]$, m by n matrices of variables over k. Let $2 \leq s_i \leq t_i$ integers, and $X_{s_1t_1}$, $Y_{s_2t_2}$ are the submatrices of X and Y coming from the first s_i rows and first t_i columns. $I = I_{s_1}(X_{s_1t_1})$, $J = I_{s_2}(X_{s_2t_2})$ the ideals of $k[X]$ generated by the maximal minors of $X_{s_1t_1}$ and the maximal minors of $X_{s_2t_2}$. Let $R_1 = k[X]/I$, $R_2 = k[X]/J$ be two determinantal rings. We consider the diagonal ideal $\mathbb{D}$ of $R_1 \otimes_k R_2$, defined via the exact sequence

$$0 \longrightarrow \mathbb{D} \longrightarrow S = R_1 \otimes_k R_2 \xrightarrow{\text{mult.}} k[X]/(I+J) \longrightarrow 0.$$

The ideal $\mathbb{D}$ is generated by the images of $x_{ij} \otimes 1 - 1 \otimes x_{ij}$ in $R_1 \otimes_k R_2$.

We write the diagonal ideal $\mathbb{D} = (\{x_{ij} - y_{ij}\})$ in

$$S = k[X_{mn}, Y_{mn}]/(I_{s_1}(X_{s_1t_1}), I_{s_2}(Y_{s_2t_2})) \cong R_1 \otimes_k R_2.$$

We have a presentation of $\mathbb{D}$,

$$S^l \xrightarrow{\phi} S^{mn} \longrightarrow \mathbb{D} \longrightarrow 0$$

From this we obtain a presentation of the symmetric algebra of $\mathbb{D}$,

$$0 \to (\text{image}(\phi)) = J \longrightarrow \text{Sym}(S^{mn}) = S[Z_{mn}] \longrightarrow \text{Sym}(\mathbb{D}) \to 0.$$

Here J is the ideal generated by the entries of the row vector $[z_{11}, z_{12}, ..., z_{1n}, ..., z_{mn}] \cdot \phi$. Hence

$$\text{Sym}(\mathbb{D}) \cong S[Z_{mn}]/J,$$

where J is generated by linear forms in the variables z_{ij}. We write $\mathcal{R}(\mathbb{D}) = S[Z_{mn}]/K$, $J \subset K$. In general K is not generated by linear forms. We can rewrite $\text{Sym}(\mathbb{D}) = S[Z_{mn}]/J = k[X_{mn}, Y_{mn}, Z_{mn}]/\mathcal{J}$ and $\mathcal{R}(\mathbb{D}) = k[X_{mn}, Y_{mn}, Z_{mn}]/\mathcal{K}$.

Theorem 4. [L] *Notation as above. Let $X_{a_1 \dots a_{s_1}}$ be the s_1 by s_1 submatrix of $X_{s_1 t_1}$ with columns $a_1, ..., a_{s_1}$, $Y_{b_1 \dots b_{s_2}}$ the s_2 by s_2 submatrix of $Y_{s_2 t_2}$ with columns $b_1, ..., b_{s_2}$, $X^{l,k}_{a_1 \dots a_{s_1}}$ the $k-l+1$ by s_1 submatrix of X with rows l, $l+1, .., k$ and columns $a_1, ..., a_{s_1}$, and similarly for Y and Z.*

We define

$$g_{ij,lk} = \begin{vmatrix} z_{ij} & z_{lk} \\ x_{ij} - y_{ij} & x_{lk} - y_{lk} \end{vmatrix}$$

$$f_{a_1, \dots, a_{s_1}} = \sum_{q=1}^{s_2} (-1)^{q+1} \left| \begin{bmatrix} Z^{q,q} \\ Y^{1,q-1} \\ X^{q+1,m} \end{bmatrix}_{a_1 \dots a_{s_1}} \right|,$$

where $1 \le a_1 < a_2 < ... < a_{s_1} \le \min(t_1, t_2)$ and $1 \le i \le m$, $1 \le l \le m$, $1 \le j \le n$, $1 \le k \le n$.

Then $\mathcal{K} = (I_{s_1}(X_{s_1 t_1}), I_{s_2}(Y_{s_2 t_2}), g_{ij,lk}, f_{a_1, \dots, a_{s_1}})$.

Definition 5. Let $1 \le a_1 < a_2 < ... < a_{s_1+k-1} \le \min(t_1, t_2)$, and $1 \le l \le k \le s_2$, we define $f^{l,k}_{a_1, \dots, a_{s_1+k-1}}$ as follow:

$$f^{l,k}_{a_1, \dots, a_{s_1+k-1}} := \sum_{r=k}^{s_2} (-1)^{r+1} \left| \begin{bmatrix} Z^{l,k-1} \\ Z^{r,r} \\ X^{1,l-1} \\ Y^{1,r-1} \\ Y^{r+1,s_1} \end{bmatrix}_{a_1, \dots, a_{s_1+k-1}} \right|$$

$$+ \sum_{r=k}^{s_2} (-1)^{r+1} \sum_{u=r+1}^{s_1} \left| \begin{bmatrix} Z^{l,k-1} \\ X^{r,r} - Y^{r,r} \\ X^{1,l-1} \\ Y^{1,r-1} \\ Y^{r+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}_{a_1, \dots, a_{s_1+k-1}} \right|.$$

Definition 6. Let $1 \leq p_1 \leq m$, $1 \leq q_1 \leq n$, $a_{s_1} < ... < a_j \leq q_1 < a_{j-1} < ... < a_1$. We define $U_{p_1,q_1,a_1,...,a_{s_1}}$ as follows:

$$U_{p_1,q_1,a_{s_1},...,a_1} := z_{p_1q_1}\left|\left[\begin{array}{cccccc} & & X^{1,p_1-1} & & & \\ x_{p_1a_{s_1}} & \cdots & x_{p_1a_j} & y_{p_1a_{j-1}} & \cdots & y_{p_1a_1} \\ & & Y^{p_1+1,s_1} & & & \end{array}\right]\right|$$

$$+\sum_{k=j+1}^{m}(x_{p_1q_1}-y_{p_1q_1})(-1)^{k+p_1}z_{p_1a_k}|X^{1,...,\hat{p_1},...,m}_{a_1,...,\hat{a_k},..a_m}|+$$

$$\sum_{u=p_1+1}^{s_1}(x_{p_1q_1}-y_{p_1q_1})\left|\left[\begin{array}{cccccc} & & X^{1,p_1-1} & & & \\ x_{p_1a_{s_1}} & \cdots & x_{p_1a_j} & y_{p_1a_{j-1}} & \cdots & y_{p_1a_1} \\ & & Y^{p_1+1,u-1} & & & \\ & & Z^{u,u} & & & \\ & & X^{u+1,s_1} & & & \end{array}\right]_{a_1,...,a_{s_1}}\right|.$$

Definition 7. Let $1 \leq b_{s_2} < ... < b_1 \leq$, $1 \leq p_1 \leq m$, $1 \leq q_1 \leq n$, $a_{s_1} < ... < a_{s_2+1} < a_{p_1} < ... < a_1$ and $a_{p_1} \leq q_1$. Let i be integer such that $1 \leq i \leq p$ and $a_{s_2+1} < b_{s_2} < ... < b_{i+1} < a_{p_1-1} \leq b_i$ and $b_l \neq a_{p_1}$ for $l \geq i+1$.

We define M_{12} as follows:

$$M_{12} = z_{p_1q_1}x_{p_1a_{p_1}}\left|\left[\begin{array}{c} X^{1,p_1-1} \\ Y^{s_2+1,s_1} \end{array}\right]_{a_1,...,a_{p_1-1},a_{s_2+1},...,a_{s_1}}\right|.$$

We define

$$W_{p_1,q_1,a_1,...,a_{p1},a_{s_2+1},...,a_{s_1},b_1,...,b_{s_2}} :=$$
$$M_{12}|Y^{1,s_1}_{b_1,...,b_{s_2}}| - \left|Y^{1,i}_{b_1,..,b_i}\right| \sum_{\{c_{i+1},...,c_{p_1},d_{p_1+1},...,d_{s_2}\}=\{b_{i+1},....,b_{s_2}\}}$$
$$\left|Y^{i+1,p_1}_{c_{i+1},..,c_{p_1}}\right| U_{p_1q_1,a_1,...,a_{p_1-1},a_{p_1},d_{p_1+1},...,d_{s_2},a_{s_2+1},...,a_{s_1}}.$$

Let $1 \leq p_1 \leq m$, $1 \leq q_1 \leq n$, $v = p_1+1,...,s_2-1$, $1 \leq a_{s_1} < ... < a_{s_2+1} < a_{p_1} < ... < a_1 \leq t_1$ and $a_{p_1} \leq q_1$. Let i be an integer, $1 \leq i \leq p$ and let $a_{s_2+1} < b_{s_2} < ... < b_{v+2} < b^{'}_v < ... < b^{'}_{p_1+1} < b_{p_1} < ... < b_{i+1} < a_{p_1-1} \leq b_{p_1+1}$ and $b^{'}_l \neq a_{p_1}$ for $l \geq i+1$ and $b^{'}_{v-1} \leq b_{v+1}$. Let $a_{s_1} < .... < a_{s_2+1} < b_{s_2} < ... < b_{v+2} < b_{v+1} < b_v < b_{v-1} < ... < b_{p_1+2} < a_{p_1} < a_{p_1-1} \leq b_{p_1+1}$, and $b^{'}_r \leq b_{r+2} < b_{r+1}$ for $r = p_1,...,v-2$.

We define

$$W^{p_1+1,v}_{p_1,q_1,a_1,...,a_{p1},a_{s_2+1},...,a_{s_1},b_1,...,b_{p_1+1},b_{p_1+2},b_{p_1+3}.,b_{s_2},b'_{p_1+1},b'_{p_1+2},...,b'_v} :=$$
$$y_{v-1,b'_{v-1}}y_{v,b_v}W^{p_1+1,v-2}_{p_1,q_1,a_1,...,a_{p1},a_{s_2+1},...,a_{s_1},b_1,...,b_{v-1},b'_v,b_{v+1},...,b_{s_2},b'_{p_1+1},b'_{p_1+2},...,b'_{v-2}}$$
$$-y_{v,b'_v}W^{p_1+1,v-1}_{p_1,q_1,a_1,...,a_{p1},a_{s_2+1},...,a_{s_1},b_1,...,b_{s_2},b'_{p_1+1},b'_{p_1+2},...,b'_{v-1}},$$

where

$$W^{p_1+1,p_1-1}_{p_1,q_1,a_1,...,a_{p_1},a_{s_2+1},...,a_{s_1},b_1,...,b_{s_2}} = U_{p_1,q_1,a_1,...,a_{s_1}}$$

and

$$W^{p_1+1,p_1}_{p_1q_1,a_1,\dots,a_{p_1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_2}} = W_{p_1q_1,a_1,..,a_{p_1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_2}}.$$

Definition 8. Let $b_{s_1} < ... < b_1$, and $1 \leq p_l < ... < p_k < b_{s_1} < ... < b_{s_2+1} < c_{s_2} < ... < c_{k+1} < b_{k-1} < ... < b_1 < a_{k-1} < ... < a_1 \leq t_1$.

Let

$$M_{12} = \left| \begin{bmatrix} Z^{l,k} \\ X^{1,k-1} \\ Y^{s_2+1,s_1} \end{bmatrix}_{p_l,..,p_k,a_1,\dots,a_{k-1},b_{s_2+1},\dots,b_{s_1}} \right|.$$

We define

$$\begin{gathered} V_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}} := \\ M_{12}|Y^{1,s_2}_{b_1,\dots,b_{s_2}}| - \sum_{\{e_k,c_{k+1},\dots,c_{s_2}\}=\{b_k,\dots.,b_{s_2}\}} \\ \pm y_{ke_k} f^{l,k}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{k-1},c_{k+1},\dots,c_{s_2},b_{s_2+1},\dots,b_{s_1}}. \end{gathered}$$

Definition 9. Let $1 \leq l \leq k \leq s_2$ and $1 \leq p_l < ... < p_k < b_{s_1} < ... < b_{s_2+1} < ... < b_{k+1} < b_{k-1} < b_k < b_{k-2} < ... < b_1 < a_{l-1} < ... < a_1 \leq t_1$. Let $w = k, ..., s_2 - 1$ and $1 \leq b_{s_2} < ... < b_{w+2} < b^{'}_w < b^{'}_{w-1} < ... < b^{'}_k < b_{k-1} < b_{k-2} ... < b_1 \leq t_2$ and $b^{'}_{w-1} \leq b_{w+1}$, and $b^{'}_r \leq b_{r+2} < b_{r+1}$ for $r = k, ..., l-2$.

We define

$$\begin{gathered} V^{k,w}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1},b'_k,b'_{k+1},\dots,b'_w} := \\ y_{w-1,b_{w-1}} y_{w,b_w} V^{k,w-2}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,..,b'_w,\dots,b_{s_1},b'_k,b'_{k+1},\dots,b'_{w-2}} \\ -y_{wb'_w} V^{k,w-1}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1},b'_k,b'_{k+1},\dots,b'_{w-1}} \qquad . \end{gathered}$$

Here $V^{k,k-2}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}} = V^{k,k-1}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}} = V_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}}$.

Definition 10. Let $1 \leq l \leq k \leq s_2$, $1 \leq q \leq n$, $1 \leq a_{s_1+k-1} < .... < a_1 \leq t_1$, $a_{s_1+k-1} < q$, $a_{j+1} \leq q < a_j$ for some $j = l-1, ..., s_1+k-3$. Let $\overline{f^{l,k,x_{l-1}}_{a_1,\dots,\hat{a_c},\dots,a_{s_1+k-1}}}$ be the determinant of matrices that coming from deleting row x_{l-1} and column a_c. We define $H^{l,k,q}_{a_1,\dots.,a_{s_1+k-1}}$ as following

$$H^{l,k,q}_{a_1,\dots,a_{s_1+k-1}} \quad = \quad z_{l-1,q} f^{l,k}_{a_1,\dots,a_{s_1+k-1}} - \sum_{c=k}^{j} (-1)^{k+c} g_{l-1,q,l-1,a_c} \overline{f^{l,k,x_{l-1}}_{a_1,\dots,\hat{a_c},\dots,a_{s_1+k-1}}}.$$

Definition 11. Let $1 \leq l \leq k \leq s_2$, $1 \leq q \leq n$, $1 \leq a_{s_1+k-1} < .... < a_1 \leq t_1$, $a_{s_1+k-1} < q$, $a_{j+1} \leq q < a_j$ for some $j = l-1, ..., s_1+k-3$. Let $a_{l+s_2-1} < b_{s_2} < ... < b_k < a_{l-1+k-1} = b_{k-1} < .... < a_{l-1+1} = b_1$.

Let

$$M = z_{l-1,q}x_{l-1,a_{j+1}} \left| \begin{bmatrix} Z^{l,k} \\ X^{1,l-2} \\ Y^{s_2+1,s_1} \end{bmatrix}_{a_{s_1+k-1},\ldots,a_{l+s_2-1},a_{l-1},\ldots,a_1} \right| .$$

We define

$$\begin{aligned} & I^{l,k,q}_{a_{s_1+k-1},\ldots,a_{l+s_2-1},a_{l-1},\ldots,a_1,b_1,\ldots,b_{s_2}} := \\ & M|Y^{1,s_2}_{b_1,\ldots,b_{s_2}}| - \sum_{\{e_k,c_{k+1},\ldots,c_{s_2}\}=\{b_k,\ldots.,b_{s_2}\}} \\ & \pm y_{ke_k} H^{l,k,q}_{a_{s_1+k-1},\ldots,a_{l+s_2-1},c_{s_2},\ldots,c_{k+1},b_{k-1},\ldots,b_1,a_{l-1},\ldots,a_1} . \end{aligned}$$

Definition 12. Let $1 \leq l \leq k \leq s_2$, $1 \leq q \leq n$, $k \leq w \leq s_2 - 1$, $1 \leq q_l < \ldots < q_k < b_{s_1} < \ldots < b_{s_2} < \ldots < b_{k+1} < b_{k-1} < b_k < b_{k-2} < \ldots < b_1 < a_{l-2} < \ldots < a_1 \leq t_1$, $q_l < q$, $q_l < a_{l-1} \leq q$. Let $w = k, \ldots, s_2 - 1$ and $1 \leq b_{s_2} < \ldots < b_{w+2} < b^{'}_w < b^{'}_{w-1} < \ldots < b^{'}_k < b_{k-1} < b_{k-2} \ldots < b_1 \leq t_2$ and $b^{'}_{w-1} \leq b_{w+1}$, and $b^{'}_r \leq b_{r+2} < b_{r+1}$ for $r = k, \ldots, l-2$.

We define

$$\begin{aligned} {}^{k,w}I^{l,k,q}_{q_l,\ldots,q_k,b_{s_1},\ldots,b_{s_2},\ldots,b_1,a_{l-1},\ldots,a_1,b'_k,\ldots,b'_w} &:= \\ y_{w-1,b'_{w-1}} y_{wb_w} {}^{k,w-2}I^{l,k,q}_{q_l,\ldots,q_k,b_{s_1},\ldots,b_{s_2},\ldots,b'_w,\ldots,b_1,a_{l-1},\ldots,a_1,b'_k,\ldots,b'_{w-2}} \\ -y_{wb'_w} {}^{k,w-1}I^{l,k,q}_{q_l,\ldots,q_k,b_{s_1},\ldots,b_{s_2},\ldots,b_1,a_{l-1},\ldots,a_1,b'_k,\ldots,b'_{w-1}} , \end{aligned}$$

where

$$\begin{aligned} {}^{k,k-2}I^{l,k,q}_{q_l,\ldots,q_k,b_{s_1},\ldots,b_{s_2},\ldots,b_1,a_{l-1},\ldots,a_1} &= {}^{k,k-1}I^{l,k,q}_{q_l,\ldots,q_k,b_{s_1},\ldots,b_{s_2},\ldots,b_1,a_{l-1},\ldots,a_1} \\ &= I^{l,k,q}_{q_l,\ldots,q_k,b_{s_1},\ldots,b_{s_2},\ldots,b_1,a_{l-1},\ldots,a_1} . \end{aligned}$$

Theorem 13. [L]*Use the notation of Definition 5, 6, 7, 7, 8, 9, 10, 11, 12 and let* $\mathcal{G} := \{|X^{1,s_1}_{a_1,..,a_{s_1}}|, |Y^{1,s_2}_{b_1,\ldots,b_{s_2}}|, g_{p_1q_1,p_2q_2}, f^{l,k}_{a_1,\ldots,a_{s_1+k-1}}, U_{p_1,q_1,a_1,\ldots,a_{s_1}}$, $W_{p,q,a_1,\ldots,a_p,a_{s_2+1},\ldots,a_{s_1},b_1,\ldots,b_{s_2}}$, $W^{p_1+1,v}_{p_1,q_1,a_1,\ldots,a_{p1},a_{s_2+1},\ldots,a_{s_1},b_1,\ldots,b_{p_1+1},b_{p_1+2},b_{p_1+3}.,b_{s_2},b'_{p_1+1},b'_{p_1+2},\ldots,b'_v}$, $V_{p_l,\ldots,p_k,a_1,\ldots,a_{k-1},b_1,\ldots,b_{s_1}}$, $V^{k,w}_{p_l,\ldots,p_k,a_1,\ldots,a_{k-1},b_1,\ldots,b_{s_1},b'_k,b'_{k+1},\ldots,b'_w}$, $H^{l,k,q}_{a_1,\ldots,a_{s_1+k-1}}$, $I^{l,k,q}_{a_{s_1+k-1},\ldots,a_{l+s_2-1},a_{l-1},\ldots,a_1,b_1,\ldots,b_{s_2}}$, ${}^{k,w}I^{l,k,q}_{q_l,\ldots,q_k,b_{s_1},\ldots,b_{s_2},\ldots,b_1,a_{l-1},\ldots,a_1,b'_k,\ldots,b'_w}\}$.

The $\mathcal{G}$ *is a Groebner basis of* $\mathcal{K}$ *with respect to the lexicographic term order and the variables ordered by* $z_{ij} > x_{lk} > y_{pq}$ *for any* i, j, l, k, p, q *and* $x_{ij} < x_{lk}$, $y_{ij} < y_{lk}$ *if* $i > l$ *or* $i = l$ *and* $j < k$ *and* $z_{ij} < z_{lk}$ *if* $i > l$ *or if* $i = l$ *and* $j > k$.

Corollary 14. *The initial ideal of*

$$\begin{aligned} \mathcal{K} \quad = \quad & (h_X + h_Y + h_g + h_f + h_U + h_W + h_{W^{p,q,l}} + h_V + \\ & h_{V^{l,k,w}} + h_{H^{l,k,q}} + h_{I^{l,k,q}} + h_{k,w I^{l,k,q}}), \end{aligned}$$

where

(1) $h_X = (\{x_{1a_1}x_{2a_2}....x_{s_1a_{s_1}} \,|\, 1 \leq a_{s_1} < a_{s_1-1} < ... < a_1 \leq t_1\}),$

(2) $h_Y = (\{y_{1a_1}y_{2a_2}...y_{s_2a_{s_2}} \,|\, 1 \leq a_{s_2} < a_{s_2-1} < ... < a_1 \leq t_2\}),$

(3) $h_g = (\{z_{ij}x_{lk} \,|\, i < l \textit{ or } i = l \textit{ and } j < k, \})$

(4) $h_f = (\{z_{lq_l}z_{l+1q_{l+1}}...z_{kq_k}x_{1a_1}...x_{l-1a_{l-1}}y_{1b_1}...y_{k-1b_{k-1}}y_{k+1b_{k+1}}...y_{s_1b_{s_1}} \,|\, 1 \leq q_l < q_{l+1} < ... < q_k < b_{s_1} < ... < b_{k+1} < b_{k-1} < ... < b_1 < a_{l-1} < ... < a_1 \leq t_1,\ 1 \leq l < k \leq s_2\}),$

(5) $h_U = (\{z_{pq}x_{1a_1}...x_{pa_p}y_{p+1a_{p+1}}...y_{s_1a_{s_1}} \,|\, p = 1, ..., s_1 - 1,\ 1 \leq q \leq n,\ 1 \leq a_{s_1} < a_{s_1-1} < ... < a_1 \leq t_1,\ a_p \leq q\}),$

(6) $h_W = (\{z_{pq}x_{1a_1}...x_{pa_p}y_{1b_1}...y_{s_1b_{s_1}} \,|\, p = 1, ..., s_1 - 1,\ 1 \leq i \leq p,\ 1 \leq b_{s_1} < ... < b_{s_2+1} < a_p < ... < a_1 \leq t_1, 1 \leq b_{s_1} < b_{s_1-1} < ... < b_{p+2} < b_p < b_{p-1} < ... < b_{i+1} < b_{p+1} < b_i < ... < b_1 \leq t_2,\ 1 \leq b_{s_1} < ... < b_{p+2} < b_p < ... < b_{i+1} < a_{p-1} < ... < a_1 \leq t_1, b_l \neq a_p, l = i+1, ..., s_1, a_{p-1} \leq b_{p+1}, a_p \leq q\}),$

(7) $h_{W^{p,q,l}} = (\{z_{pq}x_{1a_1}...x_{pa_p}y_{1b_1}...y_{p-1b_{p-1}}y_{pb_p}\ y_{p+1b_{p+1}}y_{p+1b'_{p+1}}...y_{vb_v}y_{vb'_v}\ y_{v+1b_{v+1}}...y_{s_1b_{s_1}} | 1 \leq p \leq m,\ 1 \leq q \leq n,\ v = p+1, ..., s_2 - 1,\ 1 \leq b_{s_1} < ... < b_{s_2+1} < a_p < ... < a_1 \leq t_1,\ a_p \leq q,\ b_{s_2+1} < b_{s_2} < ... < b_{v+2} < b'_v < ... < b'_{p+1} < b_{p-1} < ... < b_{i+1} < a_{p-1} \leq b_{p+1} < b_i < ... < b_1 \leq t_2,\ 1 \leq i \leq p,\ b'_l \neq a_p,\ l \geq i+1,\ b'_{v-1} \leq b_{v+1},\ b_{s_1} < .... < b_{s_2+1} < b_{s_2} < ... < b_{v+2} < b_{v+1} < b_v < b_{v-1} < ... < b_{p+2} < b_{p+1} < a_p < a_{p-1} \leq b_p,\ b'_r \leq b_{r+2} < b_{r+1},\ r = p, ..., v-2\}),$

(8) $h_V = (\{z_{lq_l}z_{l+1q_{l+1}}...z_{kq_k}x_{1a_1}...x_{l-1a_{l-1}}y_{1b_1}...y_{k-1b_{k-1}}y_{kb_k}\ y_{k+1b_{k+1}}...y_{s_1b_{s_1}} \,|\, 1 \leq l < k \leq s_2,\ 1 \leq q_l < q_{l+1} < ... < q_k < b_{s_1} < ... < b_{k+2} < b_k < b_{k-1} < ... < b_1 < a_{l-1} < ... < a_1 \leq t_1,\ b_{k-1} \leq b_{k+1} \leq t_2 - k + 1\},$

(9) $h_{V^{l,k,w}} = (\{z_{lq_l}...z_{kq_k}x_{1a_1}...x_{l-1a_{l-1}}y_{1b_1}...y_{k+1b_{k+1}}y_{k+1b'_{k+1}}...\ y_{wb_w}y_{wb'_w}\ y_{w+1b_{w+1}}...y_{s_1b_{s_1}} \,|\ 1 \leq l \leq k \leq s_2,\ 1 \leq p_l < ... < p_k < b_{s_1} < ... < b_{s_2+1} < ... < b_{k+1} < b_{k-1} < b_k < b_{k-2} < ... < b_1 < a_{l-1} < ... < a_1 \leq t_1,\ w = k, ..., s_2 - 1,\ 1 \leq b_{s_2} < ... < b_{w+2} < b'_w < b'_{w-1} < ... < b'_k < b_{k-1} < b_{k-2}... < b_1 \leq t_2,\ b'_{w-1} \leq b_{w+1},\ b'_r \leq b_{r+2} < b_{r+1},\ r = k, ..., l-2\}),$

(10) $h_{H^{l,k,q}} = (\{z_{l-1,q}z_{l,q_l}...z_{kq_k}x_{1a_1}...x_{l-2,a_{l-1}}y_{1b_1}...y_{k-1b_{k-1}}y_{k+1b_{k+1}}...\ y_{s_1b_{s_1}} \,|\, 1 \leq l \leq k \leq s_2,\ 1 \leq q \leq n,\ 1 \leq q_l < ... < q_k < b_{s_1} < ... < b_{k+1} < b_{k-1} < ... < b_1 < a_{l-2} < ... < a_1 \leq t_1,\ q_l < a_{l-1} \leq q < b_1\}),$

(11) $h_{I^{l,k,q}} = (\{z_{l-1,q}z_{l,q_l}...z_{kq_k}x_{1a_1}...x_{l-2,a_{l-1}}y_{1b_1}...y_{k-1b_{k-1}}y_{kb_k}\ y_{k+1b_{k+1}}...y_{s_1b_{s_1}} \,|\, 1 \leq l \leq k \leq s_2,\ 1 \leq q \leq n,\ 1 \leq q_l < ... < q_k < b_{s_1} < ... < b_{k+1} < b_{k-1} < b_k < b_{k-2} < ... < b_1 < a_{l-2} < ... < a_1 \leq t_1,\ q_l < a_{l-1} \leq q < b_1\}),$

(12) $h_{k,w\,I^{l,k,q}} = (\{z_{l-1,q}z_{l,q_l}...z_{kq_k}x_{1a_1}...x_{l-2,a_{l-1}}y_{1b_1}...y_{kb_k}y_{kb'_k}...y_{wb_w}y_{wb'_w}\ y_{w+1b_{w+1}}...y_{s_1b_{s_1}} \,|\, 1 \leq l \leq k \leq s_2,\ 1 \leq q \leq n,\ 1 \leq q_l < ... < q_k < b_{s_1} < ... < b_{k+1} < b_{k-1} < b_k < b_{k-2} < ... < b_1 < a_{l-2} < ... < a_1 \leq t_1,\ q_l < a_{l-1} \leq q < b_1,\ w = k, ..., s_2 - 1,\ 1 \leq b_{s_2} < ... < b_{w+2} < b'_w < b'_{w-1} < ... < b'_k < b_{k-1} < b_{k-2}... < b_1 \leq t_2,\ b'_{w-1} \leq b_{w+1},\ b'_r \leq b_{r+2} < b_{r+1},\ r = k, ..., l-2\}).$

3. ALEXANDER DUAL IDEALS

From Corollary 14, we see that the initial ideal of $\mathcal{K}$ is generated by square free monomials. We know that an ideal generated by square free monomials defines a Stanley-Reisner ring. Hence we can find the Alexander dual ideal of this ideal, [B-H]. We recall the definition of Alexander dual ideal.

Definition 15. If I is an ideal of $R = k[x_1, ..., x_n]$ generated by square-free monomials $(f_1, ..., f_l)$, then the Alexander dual ideal I^* of I is $\cap_i P_{f_i}$, where for any square-free monomial $f = x_{i_1} \cdots x_{i_r}$, $P_f = (x_{i_1}, ..., x_{i_r})$.

From Corollary 14, we see that each summand of the initial ideal has a similar structure as the ideal in the following lemma. Hence we find the Alexander dual ideal of this ideal first.

Lemma 16. *Let $R = k[X]$, where $X = [x_{ij}]$, $i = 1, ..., m$, $j = 1, ..., n$ and $m \leq n$. Let I be the ideal generated by $\{x_{1a_1}x_{2a_2}x_{3a_3}...x_{ma_m}\}$ with $1 \leq a_1 < a_2 < ... < a_l \leq a_{l+1} < ... < a_m \leq n$ for some $1 \leq l \leq m-1$. Then I^*, the Alexander dual ideal of I, is generated by*

$$\{\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+2}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \cdots \prod_{i_l=k_{l-1}+2}^{k_l} x_{li_l}$$
$$\prod_{i_{l+1}=k_l+1}^{k_{l+1}} x_{l+1 i_{l+1}} \prod_{i_{l+2}=k_{l+1}+2}^{k_{l+2}} x_{l+2 i_{l+2}} \cdots \prod_{i_m=k_{m-1}+2}^{n} x_{mi_m}\}$$

where $0 \leq k_1 < k_2 < k_3 ... < k_l \leq k_{l+1} < ... < k_{m-1} < n$.

Proof. Without lost of generality, we may assume $l = 1$. Induction on m, we consider $m = 2$ and $m = 3$ first. When $m = 2$, $I = (\{x_{1a_1}x_{2a_2} \mid 1 \leq a_1 \leq a_2 \leq n\})$. Now

$$\begin{aligned} I^* &= \cap_{1\leq a_1\leq a_2\leq n}(x_{1a_1},\, x_{2a_2}) = \cap_{1\leq a_1\leq n}(\cap_{a_1\leq a_2\leq n}(x_{1a_1},\, x_{2a_2})) \\ &= \cap_{1\leq a_1\leq n}(x_{1a_1}, \prod_{i_2=a_1}^{n} x_{2i_2}) = (\{\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{n} x_{2i_2} \mid 0 \leq k_1 \leq n\}). \end{aligned}$$

When $m = 3$, we have $I = (\{x_{1a_1}x_{2a_2}x_{3a_3} \mid 1 \leq a_1 \leq a_2 < a_3 \leq n\})$ Now

$$\begin{aligned} I^* &= \cap_{1\leq a_1\leq a_2< a_3< n}(x_{1a_1},\, x_{2a_2}, x_{3a_3}) \\ &= \cap_{1\leq a_1< n}(\cap_{a_1\leq a_2< n}(\cap_{a_2< a_3\leq n}(x_{1a_1},\, x_{2a_2}, x_{3a_3}))) \\ &= \cap_{1\leq a_1< n}(\cap_{a_1\leq a_2< n}(x_{1a_1}, x_{2a_2}, \prod_{i_3=a_2+2}^{n} x_{3i_3}) \\ &= \cap_{1\leq a_1< n}(x_{1a_1}, \{\prod_{i_2=a_1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{n} x_{3i_3} \mid a_1 - 1 \leq k_2 < n\}) \\ &= (\{\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{n} x_{3i_3} \mid 0 \leq k_1 \leq k_2 < n\}). \end{aligned}$$

When $m > 3$, we have

$$\begin{aligned}
I^* &= \cap_{1\le a_1\le a_2<a_3<\ldots<a_m\le n}(x_{1a_1},\, x_{2a_2}, \ldots,\, x_{ma_m}) \\
&= \cap_{1\le a_1\le a_2<a_3<\ldots<a_m\le n}(\cap_{a_1\le a_2<a_3<\ldots<a_m\le n}(x_{1a_1},\, x_{2a_2}, \ldots,\, x_{ma_m})) \\
&= \cap_{1\le a_1\le a_2<a_3<\ldots<a_m\le n}(x_{1a_1}, \{\prod_{i_2=a_1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \ldots \prod_{i_m=k_{m-1}+2}^{n} x_{mi_m} | \\
&\quad a_1-1\le k_2<k_3<\ldots<k_{m-1}<n\}) \\
&= (\{\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \ldots \prod_{i_m=k_{m-1}+2}^{n} x_{mi_m} | \\
&\quad 1\le k_1\le k_2<k_3<\ldots<k_{m-1}<n\}),
\end{aligned}$$

where the third equality comes from the induction. □

We obtain the Alexander dual ideal of in($\mathcal{L}$).

Lemma 17.

$$\begin{aligned}
(\mathrm{in}(\mathcal{L}))^* &= (h_X)^* \cap (h_Y)^* \cap (h_g)^* \cap (h_f)^* \cap (h_U)^* \cap (h_W)^* \cap (h_{W^{p,q,v}})^* \cap \\
&\quad (h_V)^* \cap (h_{V^{l,k,w}})^* \cap (h_H)^* \cap (h_I)^* \cap (h_{{k,w}I^{l,k,q}})^*
\end{aligned}$$

where

(1)

$$(h_X)^* = (\{\prod_{a_1=A_2+2}^{t_1} x_{1a_1} \prod_{a_2=A_3+2}^{A_2} x_{2a_2} \ldots \prod_{a_{s_1}=1}^{A_{s_1}} x_{s_1a_{s_1}} | 0\le A_{s_1}<\ldots<A_2<t_1\}),$$

(2)

$$(h_Y)^* = (\{\prod_{b_1=B_2+2}^{t_2} y_{1b_1} \prod_{b_2=B_3+2}^{B_2} y_{2b_2} \ldots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2b_{s_2}} | 0\le B_{s_2}<\ldots<B_2<t_2\}),$$

(3)

$$(h_g)^* = (\{\prod_{(i,j)=(1,1)}^{(u_1,u_2)} z_{ij} \prod_{(l,k)=(u_1,u_2)\uplus 2}^{(m,n)} x_{lk} | (0,0)\le (u_1,u_2)<(m,n)\}),$$

(4)

$$(h_f)^* = \bigcap_{l=1}^{s_2} \bigcap_{k=l}^{s_2} (\{ \prod_{q_l=1}^{Q_l} z_{l,q_l} \cdots \prod_{q_k=Q_{k-1}+2}^{Q_k} z_{kq_k} \prod_{a_1=A_2+2}^{t_1} x_{1a_1} \cdots \prod_{a_{l-1}=B_1+2}^{A_{l-1}} x_{l-1a_{l-1}}$$
$$\prod_{b_1=B_2+2}^{B_1} y_{1b_1} \cdots \prod_{b_{k-1}=B_{k+1}+2}^{B_{k-1}} y_{k-1b_{k-1}} \prod_{b_{k+1}=B_{k+2}+2}^{B_{k+1}} y_{k+1b_{k+1}} \cdots$$
$$\prod_{b_{s_1}=Q_k+2}^{B_{s_1}} y_{s_1b_{s_1}} | 0 \leq Q_l < \ldots < Q_k < B_{s_1} < \ldots < B_{k+1} < B_{k-1} < \ldots$$
$$< B_1 < A_{l-1} < \ldots < A_1 < t_1 \}),$$

(5)

$$(h_U)^* = \bigcap_{p=1}^{s_1-1} (\{ \prod_{q_p=A_p+1}^{n} z_{p,q_p} \prod_{a_1=A_2+2}^{t_1} x_{1a_1} \cdots \prod_{min(A_p,A_{p+1})+2}^{A_{p-1}} x_{p-1,a_{p-1}}$$
$$\prod_{a_p=1}^{A_p} x_{pa_p} \prod_{a_{p+1}=A_{p+2}+2}^{A_{p+1}} y_{p+1a_{p+1}} \cdots \prod_{a_{s_1}=1}^{A_{s_1}} y_{s_1a_{s_1}} | 0 \leq A_{s_1} < \ldots < A_{p+1} <$$
$$A_{p-1} < \ldots < A_2 < t_1, 0 \leq A_p < A_{p-1} \}),$$

(6)

$$(h_W)^* = \bigcap_{p=1}^{s_1-1} \bigcap_{i=1}^{p} (\{ \prod_{q=A_p+1}^{n} z_{pq} \prod_{a_1=A_2+2}^{t_1} x_{1a_1} \cdots \prod_{a_{p-1}=B_{i+1}+2}^{A_{p-1}} x_{p-1,a_{p-1}} \prod_{a_p=1}^{A_p} x_{pa_p}$$
$$\prod_{b_1=B_2}^{t_2} y_{1b_1} \cdots \prod_{b_{p-1}=B_p+2}^{B_{p-1}} y_{p-1b_{p-1}} \prod_{b_p=B_{p+2}+2}^{B_p} y_{pb_p} \prod_{b_{p+1}=A_{p-1}+1}^{B_{p+1}} y_{p+1b_{p+1}} \cdots$$
$$\prod_{1}^{B_{S_1}} y_{s_1b_{s_1}} | \, 0 \leq B_{s_1} < \ldots < B_{s_2+1} < A_p < \ldots < A_2 < t_1, \, 1 \leq B_{s_1} < B_{s_1-1}$$
$$< \ldots < B_{p+2} < B_p < B_{p-1} < \ldots < B_{i+1} < B_{p+1} < B_i < \ldots < B_2 < t_2,$$
$$1 \leq B_{s_1} < \ldots < B_{p+2} < B_p < \ldots < B_{i+1} < A_{p-1} < \ldots < A_2 < t_1 \}),$$

(7)

$$(h_{W^{p,q,v}})^* = \bigcap_{p=1}^{s_2-1} \bigcap_{v=p+1}^{s_2-1} \bigcap_{i=1}^{p} (\{ \prod_{q_p=A_p+1}^{s_1} z_{pq_p} \prod_{a_1=A_2+2}^{A_1} x_{1a_1} \cdots \prod_{a_p=B_{p+1}+2}^{A_p} x_{pa_p}$$

$$\prod_{b_1=B_2+2}^{t_2} y_{1b_1} \cdots \prod_{b_{p-1}=B'_{p+1}+2}^{B_{p-1}} y_{p-1b_{p-1}} \prod_{b_p=A_{p-1}+1}^{B_p} y_{pb_p} \prod_{b_{p+1}=A_{p-1}+1}^{B_{p+1}} y_{p+1b_{p+1}}$$

$$\prod_{b'_{p+1}=B'_{p+2}+2}^{B'_{p+1}} y_{p+1b'_{p+1}} \prod_{b_{p+2}=B_{p+3}+2}^{B_{p+2}} y_{p+2b_{p+2}} \prod_{b'_{p+2}=B'_{p+3}+2}^{B'_{p+2}} y_{p+2b'_{p+2}} \cdots$$

$$\prod_{b_v=min(B_{v+1}+2,B'_{v-2}+1)}^{B_v} y_{vb_v} \prod_{b'_v=B_{v+2}+2}^{B'_v} y_{vb'_v} \prod_{b_{v+1}=B'_{v-1}+1}^{B_{v+1}} y_{v+1b_{v+1}} \cdots$$

$$\prod_{b_{s_1}=A_{s_2+1}+2} y_{s_1b_{s_1}} | 0 \leq B_{s_1} < \ldots < B_{s_2+1} < A_p < \ldots < A_2 < t_1, B_{s_2+1} <$$

$$B_{s_2} < \ldots < B_{v+2} < B'_v < \ldots < B'_{p+1} < B_p < \ldots < B_{i+1} < A_{p-1} \leq B_{p+1}$$

$$< B_i < \ldots < B_2 < t_2, B_{s_1} < \ldots. < B_{s_2+1} < B_{s_2} < \ldots < B_{v+2} < B_{v+1} <$$

$$B_v < B_{v-1} < \ldots < B_{p+2} < A_p < A_{p-1} \leq B_{p+1}, B'_r \leq B_{r+2}, r = p, \ldots, v-2\}),$$

(8)

$$(h_V)^* = \bigcap_{l=1}^{s_1-1} \bigcap_{k=l}^{s_1} (\{ \prod_{q_l=1}^{Q_l} z_{lq_l} \cdots \prod_{q_k=Q_{k-1}+2}^{Q_k} z_{kq_k} \prod_{a_1=A_2+2}^{t_1} x_{1a_1} \cdots \prod_{a_{l-1}=B_1+2}^{A_{l-1}} x_{l-1a_{l-1}}$$

$$\prod_{b_1=B_2+2}^{B_1} y_{1b_1} \cdots \prod_{b_{k-1}=B_{k+1}+2}^{B_{k-1}} y_{k-1b_{k-1}} \prod_{b_k=B_{k-1}+2}^{B_k} y_{kb_k} \prod_{b_{k+1}=B_{k+2}+1}^{B_{k+1}} y_{k+1b_{k+1}} \cdots$$

$$\prod_{b_{s_1}=Q_k+2}^{B_{s_1}} y_{s_1b_{s_1}} | 0 \leq Q_l < Q_{l+1} < \ldots < Q_k < B_{s_1} < \ldots < B_{k+2} < B_k <$$

$$B_{k-1} < \ldots < B_1 < A_{l-1} < \ldots < A_2 < t_1,\ B_{k-1} \leq t_2 - k + 1\}),$$

(9)

$$\begin{aligned}
(h_{V^{l,k,w}})^* = \bigcap_{l=1}^{s_2-1} \bigcap_{k=l}^{s_2} \bigcap_{w=k}^{s_1} (\{ \prod_{q_l=1}^{Q_l} z_{lq_l} \cdots \prod_{q_k=Q_{k-1}+2}^{Q_k} z_{kq_k} \prod_{a_1=A_2+2}^{t_1} x_{1a_1} \cdots \\
\prod_{a_{l-1}=B_1+2}^{A_{l-1}} x_{l-1a_{l-1}} \prod_{b_1=B_2+2}^{B_1} y_{1b_1} \cdots \prod_{b_{k-1}=B_k+2}^{B_{k-1}} y_{k-1b_{k-1}} \prod_{b_k=B'_{k+1}+2}^{B_k} y_{kb_k} \\
\prod_{b_{k+1}=min(B_{k+2}+2, B_{k-1}+1)}^{t_2-k+1} y_{k+1b_{k+1}} \prod_{b'_{k+1}=B'_{k+2}+2}^{B'_{k+1}} y_{k+1b'_{k+2}} \cdots \\
\prod_{b_{w-1}=min(B'_{w-2}, B_w)+2}^{B_{w-1}} y_{w-1b_{w-1}} \prod_{b'_{w-1}=B_{w+1}+2}^{B'_{w-1}} y_{w-1b'_{w-1}} \prod_{b_w=B_{w+1}+2}^{B_w} y_{wb_w} \\
\prod_{b'_w=B_{w+2}+2}^{B'_w} y_{wb'_w} \prod_{b_{w+1}=B^2_w+2}^{B_{w+1}} y_{w+1b_{w+1}} \prod_{b_{w+2}=B_{w+3}+2}^{B_{w+2}} y_{w+2b_{w+2}} \cdots \prod_{b_{s_1}=Q_k+2}^{B_{s_1}} y_{s_1b_{s_1}} | \\
0 \leq Q_l < Q_{l+1} < ... < Q_k < B_{s_1} < ... < B_{t+2} < B^2_t < B^2_{t-1} < ... < B^2_{k+1} \\
< B_k < B_{k-1} < ... < B_1 < A_{l-1} < ... < A_2 < t_1, B^2_{w-1} \leq B_{w+1}, \\
B^2_r \leq B^1_{r+1} < B^1_r, r = k+1, ..., w-2, 0 \leq Q_{p+1} < Q_{p+2} < .... < Q_k < B_{s_1} \\
< ... < B_{k+1} < B_{k-1} < ... < B_1 < A_{p-1} < A_{p-2} < ... < A_1 < t_1 \}),
\end{aligned}$$

(10)

$$\begin{aligned}
(h_{H^{l,k,q}})^* = \bigcap_{l=1}^{s_2} \bigcap_{k=l}^{s_2} (\{ \prod_{q=A_{l-1}+1}^{Q} z_{l-1,q} \prod_{q_l=1}^{Q_l} z_{l,q_l} \cdots \prod_{q_k=Q_{k-1}+2}^{Q_k} z_{kq_k} \prod_{a_1=A_2+2}^{t_1} x_{1a_1} \cdots \\
\prod_{a_{l-1}=B_1+2}^{A_{l-1}} x_{l-2,a_{l-1}} \prod_{b_1=min(B_2,Q)+2}^{B_1} y_{1b_1} \cdots \prod_{b_{k-1}=B_{k+1}+2}^{B_{k-1}} y_{k-1b_{k-1}} \\
\prod_{b_{k+1}=B_{k+2}+2}^{B_{k+1}} y_{k+1b_{k+1}} \cdots \prod_{b_{s_1}=Q_k+2}^{B_{s_1}} y_{s_1b_{s_1}} | 1 \leq Q_l < ... < Q_k < B_{s_1} < ... < \\
B_{k+1} < B_{k-1} < ... < B_1 < A_{l-2} < ... < A_2 < t_1, \; Q_l < A_{l-1} \leq Q < \\
B_1 \}),
\end{aligned}$$

$$(h_{I^{l,k,q}})^* = \bigcap_{l=1}^{s_2} \bigcap_{k=l}^{s_2} (\{ \prod_{q=A_{l-1}+1}^{Q} z_{l-1,q} \prod_{q_l=1}^{Q_l} z_{l,q_l} \cdots \prod_{Q_k=Q_{k-1}+2}^{Q_k} z_{kq_k} \prod_{a_1=A_2+2}^{A_1} x_{1a_1} \cdots \prod_{a_{l-2}=B_1+2}^{A_{l-2}} x_{l-2,a_{l-1}} \prod_{b_1=min(B_2,Q)+2}^{B_1} y_{1b_1} \cdots \prod_{b_{k-1}=B_{k+1}+2}^{B_{k-1}} y_{k-1b_{k-1}} \prod_{b_k=B_{k-1}+2}^{B_k} y_{kb_k} \prod_{b_{k+1}=B_{k+2}+2}^{B_{k+1}} y_{k+1b_{k+1}} \cdots \prod_{b_{s_1}=Q_k+2}^{B_{s_1}} y_{s_1b_{s_1}} | 1 \leq Q_l < ... < Q_k < B_{s_1} < ... < B_{k+1} < ... < B_{k+1} < B_{k-1} < B_k < B_{k-2} < ... < B_1 < A_{l-2} < ... < A_2 < t_1, Q_l < A_{l-1} \leq Q < B_1 \}), \tag{11}$$

$$(h_{k,w\,I^{l,k,q}})^* = \bigcap_{l=1}^{s_2} \bigcap_{k=l}^{s_2} \bigcap_{w=k}^{s_2} (\{ \prod_{q=A_{l-1}+1}^{Q} z_{l-1,q} \prod_{q_l=1}^{Q_l} z_{l,q_l} \cdots \prod_{q_k=Q_{k-1}+2}^{Q_k} z_{kq_k} \prod_{a_1=A_2+2}^{t_1} x_{1a_1} \cdots \prod_{a_{l-2}=B_1+2}^{A_{l-2}} x_{l-2,a_{l-1}} \prod_{b_1=min(Q,B_2)+2}^{B_1} y_{1b_1} \cdots \prod_{b_{k-1}=B_{k+1}+2}^{B_{k-1}} y_{k-1b_{k-1}} \prod_{b_k=B_{k-1}+2}^{B_k} y_{kb_k} \prod_{b'_k=B'_{k+1}+2}^{B'_k} y_{kb'_k} \cdots \prod_{b_w=B_{w+1}+2}^{B_w} y_{wb_w} \prod_{b'_w=B_{w+2}+2}^{B'_w} y_{wb'_w} \cdots \prod_{b_{w+1}=B'_{w-1}+1}^{B_{w+1}} y_{w+1b_{w+1}} \prod_{b_{s_1}=Q_k+2}^{B_{s_1}} y_{s_1b_{s_1}} | 1 \leq Q_l < ... < Q_k < B_{s_1} < ... < B_{k+1} < B_{k-1} < B_k < B_{k-2} < ... < B_1 < A_{l-2} < ... < A_2 < t_1, Q_l < A_{l-1} \leq Q < B_1, 1 \leq B_{s_2} < ... < B_{w+2} < B'_w < B'_{w-1} < ... < B'_k < B_{k-1} < B_{k-2} ... < B_2 < t_2, B'_{w-1} \leq B_{w+1}, B'_r \leq B_{r+2} < B_{r+1}, k \leq r \leq l-2 \}). \tag{12}$$

Proof. This follows from Lemma 16. □

Having the Alexander dual ideal of $\text{in}(\mathcal{L})$, we can use Theorem 3 to show that $\text{in}(\mathcal{L})$ is Cohen-Macaulay once we show that the Alexander dual ideal has a linear free resolution. We recall the definition of a linear free solution and the regularity of an ideal.

Definition 18. (a) Let

$$\mathbf{F} : ... \longrightarrow F_i \longrightarrow F_{i-1} \longrightarrow ... \longrightarrow F_0$$

be a minimal homogeneous free resolution of an ideal I in a ring $R = k[x_1, ..., x_n]$ with $F_i = \oplus_j R(-a_{ij})$. We say I has a linear free resolution if $a_{ij} = a_i$ and $a_{i+1} = a_i + 1$.

(b) The regularity of I is defined as $\text{reg}(I) = \max_{i,j}\{a_{ij} - i\}$.

Fact 19. *If all the minimal homogeneous generators of I have the same degree, d, then I has a linear free resolution if and only if* $\mathrm{reg}(I) = d$.

We will show that $(\mathrm{in}(\mathcal{L}))^*$ is generated in the same degree d and $\mathrm{reg}(\mathrm{in}(\mathcal{L}))^* = d$. Before that we show the following result first. The reason we show this is that we will need the technique of the proof for the case $(\mathrm{in}(\mathcal{L}))^*$.

Lemma 20. *Let $R = k[X]$, where $X = [x_{ij}]$, $i = 1, ..., m$, $j = 1, ..., n$. Let I be the ideal generated by $\{x_{1a_1}x_{2a_2}x_{3a_3}...x_{ma_m}\}$ with $1 \leq a_1 < a_2 < ... < a_l \leq a_{l+1} < ... < a_m < n$ for some $1 \leq l \leq m-1$. Then I^*, the Alexander dual ideal of I has a linear free resolution.*

Proof. From Lemma 16, we see that I^* is generated by elements of degree $n-(m-2)$, denoted by $\mathrm{d}(I^*)$. Using Fact 19, it's sufficient to show that $\mathrm{reg}(I^*) = \mathrm{d}(I^*) = n-(m-2)$. We will induct on n to show that there is a linear filtration on I^*.

We write down I^* first,

$$I^* = (\{\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+2}^{k_2} x_{2i_2} ... \prod_{i_l=k_{l-1}+2}^{k_l} x_{ii_l} \prod_{i_{l+1}=k_l+1}^{k_{l+1}} x_{ii_{l+1}} ... \prod_{i_m=k_{m-1}+2}^{n} x_{mi_m}\})$$

with $0 \leq k_1 < k_2 < ... < k_l \leq k_{l+1} < ... < k_{m-1} < n$. When $m = n$, we need $0 \leq k_1 < k_2 < ... < k_l \leq k_{l+1} < ... < k_{m-1} < m$. Without lost of generality, we assume $l = 1$. Hence

$$I^* = (x_{11}x_{12},\ \{x_{11}x_{ii} \,|\, i = 2, ..., m\},\ \{x_{i,i-1}x_{j,j} \,|\, i = 2, ..., m,\ j = i, ..., m\}).$$

Now look at $x_{m,m}$, the terms $x_{i,i-1}x_{mm}$ for $i = 2, ..., m-1$ and $x_{11}x_{m,m}$ are multiple of $x_{m,m}$. Also $x_{i,i-1}x_{j,j}$ for $i = 2, ..., m-1$, $j =, ..., m-1$ is divisible by $x_{i,i-1}$ and $x_{11}x_{12}$ and $x_{11}x_{ii}$ for $i = 2, ..., m-1$ is divisible by x_{11}. We can rewrite

$$I^* = (\{x_{i,i-1}(x_{i,i}, ..., x_{m,m}) \,|\, i = 2, ..., m-1\},\ x_{11}(x_{12}, x_{2,2}, ..., x_{m,m})).$$

Then we have

$$\begin{aligned}
I^* = J_1 \quad &\subset \quad (\{x_{i,i-1}(x_{i,i}, ..., x_{m,m}) \,|\, i = 2, ..., m-1\}, x_{11}) = J_2 \\
&\subset \quad (\{x_{i,i-1}(x_{i,i}, ..., x_{m,m}) \,|\, i = 3, ..., m-1\}, x_{11}, x_{21}) = J_3 \\
&\subset \quad (\{x_{i,i-1}(x_{i,i}, ..., x_{m,m}) \,|\, i = 4, ..., m-1\}, x_{11}, x_{21}, x_{32}) = J_4 \\
&\subset \quad ... \\
&\subset \quad (\{x_{i,i-1} \,|\, i = 2, ..., m-1\}, x_{11}) = J_m.
\end{aligned}$$

J_m is generated by a regular sequence of degree 1, hence it has $\mathrm{reg}J_m$=1.

We will show $\mathrm{reg}(J_{l+1}/J_l) = 1$ for $l = 1, ..., m-1$. Then $\mathrm{reg}(J_l) = 2 = m-(m-2)$ for $l = 1, ..., m-1$. We write down

$$J_{l+1} = (\{x_{i,i-1}(x_{i,i}, ..., x_{m,m}) \,|\, i = l+1, ..., m-1\}, x_{11}, x_{21}, x_{32}, ..., x_{l-1,l-2}, x_{l,l-1})$$

and

$$\begin{aligned}
J_l &= (\{x_{i,i-1}(x_{i,i}, ..., x_{m,m}) \,|\, i = l, ..., m-1\}, x_{11}, x_{21}, x_{32}, ..., x_{l-1,l-2}) \\
&= (\{x_{i,i-1}(x_{i,i}, ..., x_{m,m}) \,|\, i = l+1, ..., m-1\},\ x_{l,l-1}(x_{l,l}, ..., x_{mm}), \\
&\quad\ x_{11}, x_{21}, x_{32}, ..., x_{l-1,l-2}).
\end{aligned}$$

Then

$$\begin{aligned} J_{l+1}/J_l &= (x_{l,l-1})/(x_{l,l-1} \cap (\{x_{i,i-1}(x_{i,i}, ..., x_{m,m})|\, i = l+1, ..., m-1\}), \\ &\quad x_{l,l-1} \cap (x_{11}, x_{21}, x_{32}, ..., x_{l-1,l-2}), x_{l,l-1}(x_{l,l}, ..., x_{mm})) \\ &= (x_{l,l-1})/x_{l,l-1}(x_{l,l}, ..., x_{mm}, x_{11}, x_{21}, x_{32}, ..., x_{l-1,l-2}). \end{aligned}$$

Since

$$\begin{aligned} &\operatorname{reg}((x_{l,l-1})/x_{l,l-1}(x_{l,l}, ..., x_{mm}, x_{11}, x_{21}, x_{32}, ..., x_{l-1,l-2})) \\ = \quad &\operatorname{reg}(R/(x_{l,l}, ..., x_{mm}, x_{11}, x_{21}, x_{32}, ..., x_{l-1,l-2})) + 1 \quad = 1, \end{aligned}$$

we have $\operatorname{reg}(J_{l+1}/J_l) = 1$ for all $l = 1, ..., m-1$.

For the induction part, we write $I^* := I^*_n$ when X is a m by n matrix. We assume $\operatorname{reg}(I^*_{n-1}) = n-1-(m-2)$ and the degree of the generating of I^*_{n-1} is $n-1-(m-2)$. We look at the variable $x_{m,n}$. When $k_{m-1} < n-1$,

$$\{\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \cdots \prod_{i_m=k_{m-1}+2}^{n} x_{m,i_m}\}$$

is divisible by $x_{m,n}$ and we write

$$\begin{aligned} A = \quad & \{\textstyle\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \cdots \prod_{i_m=k_{m-1}+2}^{n} x_{m,i_m}| \\ & \quad 0 \le k_1 \le k_2 < k_3 < ... < k_{m-1} = n-2\} \\ = \quad & \{(\textstyle\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \cdots \prod_{i_m=k_{m-2}+2}^{n-2} x_{m-1,i_{m-1}})x_{m,n}| \\ & \quad 0 \le k_1 \le k_2 < k_3 < ... < k_{m-1} = n-2\} \\ = \quad & A' x_{m,n} \end{aligned}$$

When $k_{m-1} = n-1$, $\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \cdots \prod_{i_m=k_{m-1}+2}^{n} x_{mi_m}$ is not divisible by $x_{m,n}$, we have

$$\begin{aligned} B = \quad & \{\textstyle\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \cdots \prod_{i_m=k_{m-1}+2}^{n} x_{mi_m}| \\ & \quad 0 \le k_1 \le k_2 < k_3 < ... < k_{m-1} = n-1\} \\ = \quad & \{\textstyle\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \cdots \prod_{i_m=k_{m-2}+2}^{n-1} x_{m-1,i_{m-1}}| \\ & \quad 0 \le k_1 \le k_2 < k_3 < ... < k_{m-2} < k_{m-1} = n-1\} \\ = \quad & \{(\textstyle\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \cdots \prod_{i_m=k_{m-2}+2}^{n-1} x_{m-1,i_{m-1}}) \\ & \quad x_{j,j-1+(n-1-(m-2))-1}|\, 0 \le k_1 \le k_2 < k_3 < ... < k_{m-2} < n-2,\ 1 < j < m\}, \\ & \{(\textstyle\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \cdots \prod_{i_m=k_{m-2}+2}^{n-1} x_{m-1,i_{m-1}}) \\ & \quad x_{1,1+(n-1-(m-2))-1}|\, 0 \le k_1 \le k_2 < k_3 < ... < k_{m-2} < n-2\}. \\ = \quad & A'(\{x_{j,j-1+(n-1-(m-2))-1}|\, 1 < j < m\}, x_{1,n-1-(m-2)}). \end{aligned}$$

When $k_{m-1} \leq n-3$, we write

$$\begin{aligned}
C &= \{\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \dots \prod_{i_m=k_{m-1}+2}^{n} x_{m,i_m} | \\
&\quad 0 \leq k_1 \leq k_2 < k_3 < \dots < k_{m-2} < k_{m-1} \leq n-3\} \\
&= \{(\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \dots \prod_{i_m=k_{m-1}+2}^{n-1} x_{m,i_m}) x_{m,n} | \\
&\quad 0 \leq k_1 \leq k_2 < k_3 < \dots < k_{m-2} < k_{m-1} \leq n-3\} \\
&= C' x_{mn}.
\end{aligned}$$

On the other hand C can be written as

$$\begin{aligned}
C &= \{(\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \dots \prod_{i_m=k_{m-1}+2}^{n-1} x_{m,i_m}) x_{j,j-1+(n-1-(m-2))-1} x_{m,n} | \\
&\quad 0 \leq k_1 \leq k_2 < k_3 < \dots < k_{m-2} < k_{m-1} - 1 < k_m \leq n-3,\ 1 < j < m\},
\end{aligned}$$

$$\begin{aligned}
&\quad \{(\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \dots \prod_{i_m=k_{m-1}+2}^{n-1} x_{m,i_m}) x_{1,1+(n-1-(m-2))-1} x_{m,n} | \\
&\quad 0 \leq k_1 \leq k_2 < k_3 < \dots < k_{m-2} < k_{m-1} - 1 < k_m \leq n-3\} \\
&= C'' x_{m,n} (\{x_{j,j-1+(n-1-(m-2))-1} | \, 1 < j < m\}, x_{1,n-1-(m-2)}).
\end{aligned}$$

Notice $I^*_{n-1} = (A^{'}, C')$ and $\mathrm{d}(A') = \mathrm{d}(C')$. We also have

$$\begin{aligned}
A' &= (\{(\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \dots \prod_{i_m=k_{m-2}+2}^{n-2} x_{m-1,i_{m-1}}) | \\
&\quad 0 \leq k_1 \leq k_2 < k_3 < \dots < k_{m-2} < k_{m-1} = n-2\}) \\
&= (\{(\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \dots \prod_{i_m=k_{m-2}+2}^{n-2} x_{m-1,i_{m-1}}) x_{j,j-1+(n-2-(m-2))-1} | \\
&\quad 0 \leq k_1 \leq k_2 < k_3 < \dots < k_{m-2} < k_{m-1} = n-3,\ 1 < j < m\}, \\
&\quad \{(\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \dots \prod_{i_m=k_{m-2}+2}^{n-2} x_{m-1,i_{m-1}}) x_{1,1+(n-2-(m-2))-1} | \\
&\quad 0 \leq k_1 \leq k_2 < k_3 < \dots < k_{m-2} < k_{m-1} = n-3\}) \\
&= A'' (\{x_{j,j-1+n-m-1} | \, 1 < j < m\}, x_{1,n-2-(m-2)}).
\end{aligned}$$

Now look at C', we obtain

$$\begin{aligned}
C' &= (\{\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \ldots \prod_{i_m=k_{m-1}+2}^{n-1} x_{m,i_m} | \\
&\quad 0 \le k_1 \le k_2 < k_3 < \ldots < k_{m-2} < k_{m-1} < n-3\}, \\
&\quad \{\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \ldots \prod_{i_m=k_{m-1}+2}^{n-1} x_{m,i_m} | \\
&\quad 0 \le k_1 \le k_2 < k_3 < \ldots < k_{m-2} < k_{m-1} = n-3\}) \\
&= (\{(\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \ldots \prod_{i_m=k_{m-1}+2}^{n-2} x_{m,i_m}) x_{m,n-1} | \\
&\quad 0 \le k_1 \le k_2 < k_3 < \ldots < k_{m-2} < k_{m-1} < n-3\}, \\
&\quad A'' x_{m,n-1}).
\end{aligned}$$

Hence

$$\begin{aligned}
(A') \cap (C') &= (A''(\{x_{j,j-1+n-m-1} | \, 1 < j < m\}, x_{1,n-2-(m-2)})) \cap \\
&\quad (\{(\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \ldots \prod_{i_m=k_{m-1}+2}^{n-2} x_{m,i_m}) x_{m,n-1} | \\
&\quad 0 \le k_1 \le k_2 < k_3 < \ldots < k_{m-2} < k_{m-1} < n-3\}, \\
&\quad A'' x_{m,n-1}) \\
&= (A''(\{x_{j,j-1+n-m-1} | \, 1 < j < m\}, x_{1,n-2-(m-2)})) \cap \\
&\quad (\{(\prod_{i_1=1}^{k_1} x_{1i_1} \prod_{i_2=k_1+1}^{k_2} x_{2i_2} \prod_{i_3=k_2+2}^{k_3} x_{3i_3} \ldots \prod_{i_m=k_{m-1}+2}^{n-2} x_{m,i_m}) x_{m,n-1} | \\
&\quad 0 \le k_1 \le k_2 < k_3 < \ldots < k_{m-2} < k_{m-1} < n-3\}), \\
&\quad (A''(\{x_{j,j-1+n-m-1} | \, 1 < j < m\}, x_{1,n-2-(m-2)})) \cap \\
&\quad (A'' x_{m,n-1}) \\
&= (A' x_{m,n-1}) \\
&= (C'(\{x_{j,j-1+n-m-1} | \, 1 < j < m\}, x_{1,n-2-(m-2)})).
\end{aligned}$$

On the other hand, we have

$$\begin{aligned}
(A') \cap (C' x_{m,n}) &= (A') \cap (C'' x_{m,n}(\{x_{j,j-1+n-m} | \, 1 < j < m\}, x_{1,n-1-(m-2)})) \\
&\subset (A'(\{x_{j,j-1+n-m} | \, 1 < j < m\}, x_{1,n-m+1})).
\end{aligned}$$

We look at the filtration:

$$\begin{aligned}
I_n^* &= (C, B, A) \\
&= (C' x_{m,n}, \ A'(\{x_{j,j-1+n-m} | \, 1 < j < m\}, x_{1,n-m+1}), \ A' x_{m,n}) \\
&\subset (C' x_{m,n}, A') \\
&\subset (C', A') = I_{n-1}^*.
\end{aligned}$$

We have

$$\begin{array}{rc} & \operatorname{reg}((C', A')/(C'x_{m,n}, A')) \\ = & \operatorname{reg}((C')/((C') \cap (A'), C'x_{m,n})) \\ = & \operatorname{reg}((C')/(C'(\{x_{j,j-1+n-m-1} \mid 1 < j < m\}, x_{1,n-2-(m-2)}, x_{m,n})) \\ = & \operatorname{reg}(R/(\{x_{j,j-1+n-m-1} \mid 1 < j < m\}, x_{1,n-2-(m-2)}, x_{mn}) + \deg(C') \\ = & \deg(C') \\ = & n-(m-2)-1. \end{array}$$

By induction hypothesis, we have $\operatorname{reg}(C', A') = \operatorname{reg}(I^*_{n-1}) = n-1-(m-2)$, hence $\operatorname{reg}(C'x_{m,n}, A') = n-(m-2)$. Similarly

$$\begin{array}{rc} & \operatorname{reg}((C'x_{m,n}, A')/(C'x_{m,n},\ A'(\{x_{j,j-1+n-m} \mid 1 < j < m\}, x_{1,n-m+1}),\ A'x_{m,n})) \\ = & \operatorname{reg}((A')/((A') \cap (C'x_{m,n}), A'(\{x_{j,j-1+n-m} \mid 1 < j < m\}, x_{1,n-m+1}),\ A'x_{m,n})) \\ = & \operatorname{reg}((A')/(A'(\{x_{j,j-1+n-m} \mid 1 < j < m\}, x_{1,n-m+1},\ x_{m,n})) \\ = & \operatorname{reg}(R/(A'(\{x_{j,j-1+n-m} \mid 1 < j < m\}, x_{1,nm+1},\ x_{m,n}))) + \mathrm{d}(A') \\ = & \mathrm{d}(A') \\ = & n-(m-2)-1. \end{array}$$

Hence $\operatorname{reg}(I^*_n) = n-(m-2)$. ☐

Lemma 21. *The Alexander dual of* $\operatorname{in}(\mathcal{L})$, $(\operatorname{in}(\mathcal{L}))^*$, *is generated by square free monomials with degree* $mn-1+t_2-(s_2-1)+t_1-(s_1-1)$ *and* $\operatorname{reg}(\operatorname{in}(\mathcal{L}))^* = mn-1+t_2-(s_2-1)+t_1-(s_1-1)$.

Proof. We prove this lemma by inducting on n. Since $s_1 > s_2$, the generators of the ideal $(\operatorname{in}(\mathcal{L}))^*$ do not involving variables x_{ij}, y_{lk}, z_{pq} when $i, l, p > s_1$. Hence we may assume $s_1 = m$. When $m = n$, we have $m = s_1 \leq t_1 \leq m$. Then we have

$$(\operatorname{in}(\mathcal{L}))^* = (h_X)^* \cap (h_Y)^* \cap (h_g)^* \cap (h_f)^* \cap (h_U)^* \cap (h_W)^* \cap (h_{W^{p,q,v}})^*.$$

We write down each component,

$$(h_X)^* = (x_{1m},\ x_{2,m-1}, ...,\ x_{m,1}),$$

$$(h_f)^* = (t_{1,1}, y_{2,m}, y_{3,m-1}, ..., y_{m2}),$$

$$(h_Y)^* \quad = \quad (\{\prod_{b_1=B_2+2}^{t_2} y_{1b_1} \prod_{b_2=B_3+2}^{B_2} y_{2b_2} \cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2 b_{s_2}} | 0 \leq B_{s_2} < ... < B_2 < t_2\}),$$

$$(h_g)^* = (\{\prod_{(i,j)=(1,1)}^{(u_1,u_2)} z_{ij} \prod_{(l,k)=(u_1,u_2)\uplus 2}^{(m,n)} x_{lk} \mid (0,0) \leq (u_1, u_2) < (m,n)\}),$$

$$\begin{aligned}
(h_U)^* \quad = \quad & \bigcap_{p=1}^{m-1} \left(\bigcap_{(p,u_2)=(p,m)}^{(p,m-(p-1))} (z_{p,u_2}, x_{1,m}, ..., x_{p,m-(p-1)}, y_{p+1,m-p}, ..., y_{m,1}) \right. \\
& \bigcap_{(p,u_2)=(p,m-(p-1)-1)}^{(p,1)} (z_{p,u_2}, x_{1,m}, ..., x_{p,u_2}, y_{p+1,m-p+1}, ..., y_{j,u_2+1}, \\
& \left. y_{j+1,u_2-1}, ..., y_{m,1}) \right),
\end{aligned}$$

$$\begin{aligned}
(h_W)^* \quad = \quad & \bigcap_{p=1}^{m-1} \bigcap_{(p,u_2)=(p,m)}^{(p,m-(p-1))} (z_{p,u_2}, x_{1,m}, ..., x_{p,m-(p-1)}, y_{1,m}, ..., y_{p-1,m-p+2}, \\
& y_{p,m-p}, y_{p+1,m-p+1}, y_{p+2,m-p-1}, ..., y_{m,1}),
\end{aligned}$$

$$\begin{aligned}
(h_{W^{p,q,v}})^* \quad = \quad & \bigcap_{p=1}^{m-1} \bigcap_{v=p+1}^{m-1} \bigcap_{(p,u_2)=(p,m)}^{(p,m-(p-1))} (z_{p,u_2}, x_{1,m}, ..., x_{p,m-(p-1)}, y_{1,m}, ..., y_{p-1,m-p+2}, \\
& y_{p.m-p}, y_{p+1,m-p+1}, y_{p+1,m-p-1}, y_{p+2,m-p}, y_{p+2,m-p-2}, ..., \\
& y_{v,m-(v-1)-1}, y_{v,m-(v-1)+1}, y_{v+1,m-(v-1)}, ..., y_{m,1}).
\end{aligned}$$

Then we have

$$\begin{aligned}
(\mathrm{in}(\mathcal{L}))^* \quad = \quad & (\{ \prod_{(i,j)=(1,1)}^{(u_1,u_2)} z_{ij} \prod_{(l,k)=(u_1,u_2)\uplus 2}^{(m,n)} x_{lk} x_{i,m-(i-1)} \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2 b_{s_2}} | \\
& (u_1, u_2) < (m-1, m-1), 1 \le i \le m, 0 \le B_{s_2} < ... < B_2 < t_2 \},
\end{aligned}$$

$$\begin{aligned}
& \{ \prod_{(i,j)=(1,1)}^{(u_1,u_2)} z_{ij} \prod_{(l,k)=(u_1,u_2)\uplus 2}^{(m,n)} x_{lk} x_{i,m-(i-1)} \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2 b_{s_2}} | \\
& (m-1, m-1) \le (u_1, u_2) < (m-2, m), 1 \le i \le m-1, 0 \le B_{s_2} < ... < B_2 < t_2 \},
\end{aligned}$$

$$\begin{aligned}
& \{ \prod_{(i,j)=(1,1)}^{(u_1,u_2)} z_{ij} \prod_{(l,k)=(u_1,u_2)\uplus 2}^{(m,n)} x_{lk} \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2 b_{s_2}} y_{m,1} | \\
& (m-1, m-1) \le (u_1, u_2) < (m-2, m), s_2 < m, 0 \le B_{s_2} < ... < B_2 < t_2 \},
\end{aligned}$$

$$\begin{aligned}
& \{ \prod_{(i,j)=(1,1)}^{(u_1,u_2)} z_{ij} \prod_{(l,k)=(u_1,u_2)\uplus 2}^{(m,n)} x_{lk} \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2 b_{s_2}} y_{m,2} | \\
& (m-1, m-1) \le (u_1, u_2) < (m-2, m), s_2 = m, 0 \le B_{s_2} < ... < B_2 < t_2 \},
\end{aligned}$$

$$\begin{aligned}
& \{ \prod_{(i,j)=(1,1)}^{(u_1,u_2)} z_{ij} \prod_{(l,k)=(u_1,u_2)\uplus 2}^{(m,n)} x_{lk} x_{i,m-(i-1)} \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2 b_{s_2}} | \\
& (m-2, m) \le (u_1, u_2) < (m-2, 1), 1 \le i \le m-2, 0 \le B_{s_2} < ... < B_2 < t_2 \},
\end{aligned}$$

$$\{\prod_{(i,j)=(1,1)}^{(m-2,m)} z_{ij} \prod_{(l,k)=(m-1,2)}^{(m,n)} x_{lk}x_{m-1,1} \prod_{b_1=B_2+2}^{t_2} y_{1b_1}\cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2b_{s_2}} | \\ 0 \leq B_{s_2} < \ldots < B_2 < t_2\},$$

$$\{\prod_{(i,j)=(1,1)}^{(m-2,m)} z_{ij} \prod_{(l,k)=(m-1,2)}^{(m,n)} x_{lk} \prod_{b_1=B_2+2}^{t_2} y_{1b_1}\cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2b_{s_2}} y_{m,2} | \\ 0 \leq B_{s_2} < \ldots < B_2 < t_2\},$$

$$\{\prod_{(i,j)=(1,1)}^{(u_1,u_2)} z_{ij} \prod_{(l,k)=(u_1,u_2)\uplus 2}^{(m,n)} x_{lk} \prod_{b_1=B_2+2}^{t_2} y_{1b_1}\cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2b_{s_2}}(y_{m,1},\ y_{m-1,2}) | \\ (m-2,m) < (u_1,u_2) < (m-2,1), 0 \leq B_{s_2} < \ldots < B_2 < t_2,\ s_2 < m\},$$

$$\{\prod_{(i,j)=(1,1)}^{(u_1,u_2)} z_{ij} \prod_{(l,k)=(u_1,u_2)\uplus 2}^{(m,n)} x_{lk} \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \prod_{b_2=B_3+2}^{B_2} y_{2b_2}\cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2b_{s_2}}(y_{m,2},\ y_{m-1,1}) | \\ (m-2,m) < (u_1,u_2) < (m-2,1),\ 0 \leq B_{s_2} < \ldots < B_2 < t_2,\ s_2 = m \text{ and} \\ \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \prod_{b_2=B_3+2}^{B_2} y_{2b_2}\cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2b_{s_2}} \text{ is divisible by } y_{m,1} \text{ or } y_{m-1,2}\},$$

$$\{\prod_{(i,j)=(1,1)}^{(u_1,u_2)} z_{ij} \prod_{(l,k)=(u_1,u_2)\uplus 2}^{(m,n)} x_{lk}x_{i,m-(i-1)} \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \prod_{b_2=B_3+2}^{B_2} y_{2b_2}\cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2b_{s_2}} | \\ (m-2,1) \leq (u_1,u_2) < (m-3,2), 1 \leq i \leq m-3,\ 0 \leq B_{s_2} < \ldots < B_2 < t_2\},$$

$$\{\prod_{(i,j)=(1,1)}^{(m-2,1)} z_{ij} \prod_{(l,k)=(m-2,3)}^{(m,n)} x_{lk}x_{m-2,2} \prod_{b_1=B_2+2}^{t_2} y_{1b_1}\cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2b_{s_2}} | \\ 0 \leq B_{s_2} < \ldots < B_2 < t_2\},$$

$$\{\prod_{(i,j)=(1,1)}^{(m-2,1)} z_{ij} \prod_{(l,k)=(m-2,3)}^{(m,n)} x_{lk} \prod_{b_1=B_2+2}^{t_2} y_{1b_1}\cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2b_{s_2}}(y_{m-1,3}, y_{m,1}) | \\ 0 \leq B_{s_2} < \ldots < B_2 < t_2,\ s_2 < m\},$$

$$\{\prod_{(i,j)=(1,1)}^{(m-2,1)} z_{ij} \prod_{(l,k)=(m-2,3)}^{(m,n)} x_{lk} \prod_{b_1=B_2+2}^{t_2} y_{1b_1}\cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2b_{s_2}}(y_{m-1,3}, y_{m,2}) | \\ 0 \leq B_{s_2} < \ldots < B_2 < t_2 \text{ and } s_2 = m, \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \prod_{b_2=B_3+2}^{B_2} y_{2b_2}\cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2b_{s_2}} \\ \text{is divisible by } y_{m,1}\},$$

$$\{ \prod_{(i,j)=(1,1)}^{(m-3,m)} z_{ij} \prod_{(l,k)=(m-2,2)}^{(m,n)} x_{lk} x_{m-2,1} \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2 b_{s_2}} | \\ 0 \leq B_{s_2} < ... < B_2 < t_2\},$$

$$\{ \prod_{(i,j)=(1,1)}^{(m-3,m)} z_{ij} \prod_{(l,k)=(m-2,2)}^{(m,n)} x_{lk} \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2 b_{s_2}} (y_{m-1,3}, y_{m,2}) | \\ 0 \leq B_{s_2} < ... < B_2 < t_2\},$$

$$\{ \prod_{(i,j)=(1,1)}^{(u_1,u_2)} z_{ij} \prod_{(l,k)=(u_1,u_2)\uplus 2}^{(m,n)} x_{lk} \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2 b_{s_2}} (y_{m,1}, y_{m-1,2}, y_{m-2,3}) | \\ (m-3, m-1) \leq (u_1, u_2) < (m-3, 2), 0 \leq B_{s_2} < ... < B_2 < t_2\}, \\ ... $$

$$\{ \prod_{(l,k)=(1,2)}^{(m,n)} x_{lk} \prod_{b_1=B_2+2}^{t_2} y_{1b_1} \prod_{b_2=B_3+2}^{B_2} y_{2b_2} \cdots \prod_{b_{s_2}=1}^{B_{s_2}} y_{s_2 b_{s_2}} (y_{2,m}, y_{3,m-1}, ..., y_{m,2}) | \\ 0 \leq B_{s_2} < ... < B_2 < t_2\}).$$

Notice that all the elements are in the same degree $mm-1+1+t_2-(s_2-1)$. We find a filtration starting from $(\text{in}(\mathcal{L}))^*$ and ending at $(x_{m,1}, x_{m-1,2}, ..., x_{1m})$. Each quotient of this filtration will have the form P/PL, where P is an ideal generated in the same degree and L is an ideal generated by variables such that those variables form a regular sequence modulo P.

We look at the variable $y_{s_2,1}$. The elements in $(\text{in}(\mathcal{L}))^*$ that are divisible by $y_{s_2,1}$ must have $B_{s_2} > 0$. The elements in $(\text{in}(\mathcal{L}))^*$ that are not divisible by $y_{s_2,1}$ must have $B_{s_2} = 0$. Hence $(\text{in}(\mathcal{L})^*) = (A_1(y_{s_2,1}, ..., y_{s_2-1,2}, ..y_{1,s_2})), C_1 y_{s_2,1})$, where the elements in C_1 have $B_{s_2} > 1$. Also all the elements of A_1 and C_1 are in the same degree. Furthermore, $A_1 \cap C_1 = A_1 c_1 = C_1(\{a_{1i}\})$, $(y_{s_2,1}, ..., y_{1,s_2})$ is a regular sequence modulo A_1 and $(y_{s_2,1}, \{a_{1i}\})$ is a regular sequence modulo C_1. We look at the following filtration

$$(\text{in}(\mathcal{L}))^* \subset (A_1, C_1 y_{s_2,1}) \subset (A_1, C_1).$$

Then we can use the proof of Lemma 20 to show that the quotients are $A_1/A_1(y_{s_2,1}, ..., y_{1,s_2})$ and $C_1/C_1(\{a_{1i}\},\ y_{s_2,1})$. Notice the following equalities: $\text{reg}(A_1/A_1(y_{s_2,1}, ..., y_{1,s_2})) = \text{reg}(A_1)$ and $\text{reg}(C_1/C_1(\{a_{1i}\}, y_{s_2,1}) = \text{reg}(C_1)$.

Next, we look at $y_{s_2,2}$ and write $(A_1, C_1) = (A_2(y_{s_2,2}, ..., y_{1,s_2+1}), C_2 y_{s_2,2})$. As before we have a filtration

$$(A_1, C_1) \subset (A_2, C_2 y_{s_2,2}) \subset (A_2, C_2),$$

and the quotients are $A_2/A_2(y_{s_2,2}, ..., y_{1,s_2+1})$ and $C_2/C_2(\{a_{2i}\}, y_{s_2,2})$. We can continue to $y_{s_2,3}$ until $y_{s_2,t_2-(s_2-1)}$, we will reduce to an ideal J_1 which is generated in the same degree $mm-1+1$.

Now we look at the variable $z_{1,1}$. When $(u_1, u_2) = (0, 0)$ in an element, $z_{1,1}$ is not a factor of this element. When $(u_1, u_2) \leq (1, 1)$, then $z_{1,1}$ is a factor. The ideal J_1 can be written as $(D_{1,1}(z_{1,1}, \{d_{1,1,i}\}),\ E_{1,1} z_{1,1})$. Hence we have a filtration

$$J_1 \subset (D_{1,1}, E_{1,1} z_{1,1}) \subset (D_{1,1}, E_{1,1})$$

with quotients $D_{1,1}/D_{1,1}(z_{1,1},\{d_{1,1,i}\})$ and $E_{1,1}/E_{1,1}(\{e_{1,1,i}\},z_{1,1})$. We look at $z_{1,2}$ next and reduce to an ideal $(D_{1,2},E_{1,2})$, we continue to $z_{m,m-1}$. We can find a filtration from J_1 to $(D_{m,m-1},E_{m,m-1})=(x_{m,1},....,x_{1m})$.

Since $\operatorname{reg}(x_{m,1},....,x_{1,m})=\operatorname{reg}(D_{m,m-1},E_{m,m-1})=1$, it follows that $\operatorname{reg}(D_{m,m-1})$ and $\operatorname{reg}(E_{m,m-1})$ are equal to 1. Also the final quotient

$$(D_{m,m-1},E_{m,m-1})/(D_{m,m-1},E_{m,m-1}z_{m,m-1})$$

has regularity equals to the regularity of $E_{m,m-1}$. It follows that

$$\operatorname{reg}(D_{m,m-1},E_{m,m-1}z_{m,m-1})=2.$$

Since the quotient $(D_{m,m-1},E_{m,m-1}z_{m,m-1})/(D_{m,m-2},E_{m,m-2})$ has regularity equal to $\operatorname{reg}(D_{m,m-1})=1$, we have $\operatorname{reg}(D_{m,m-2},E_{m,m-2})=2$. Continue with the same argument, we obtain $\operatorname{reg}(J_1)=1+mm-1$. Hence $\operatorname{reg}((\operatorname{in}\mathcal{L})^*)=1+mm-1+t_2-(s_2-1)$.

For the induction steps, we write I_n for the Alexander dual ideal $(\operatorname{in}(\mathcal{L}))^*$ in the m by n matrix case. We assume by induction hypothesis that $\operatorname{reg}(I_{n-1})=\operatorname{d}(I_{n-1})=m(n-1)-1+t_1-(s_1-1)+t_2-(s_2-1)$. We will show that we can build a filtration from I_n to I_{n-1} such that the quotients are the form P/PL where P is an ideal and L is an ideal generated by variables such that they are a regular sequence modulo P. From Lemma 17 we know that each summand of $\operatorname{in}(\mathcal{L})$ is generated by monomials satisfying the assumption of Lemma 16. For a fixed variable y_{ij}, we can write $(h_Y)^*=(A^Yy_{ij},A^Y(\{a_l^Y\}),B^Yy_{ij})$. Also $A^Y\cap B^Y=B^Y(\{b_i^Y\})$, where $\{b_i^Y\}$ are variables such that $(y_{ij},\{b_i^Y\})$ is a regular sequence modulo B^Y and $(y_{ij},\{y_{lk}\})$ is a regular sequence modulo A^Y. Similarly, $(h_U)^*=(A^Uy_{ij},A^U(\{y_{lk},x_{p,q},z_{uv}\}),B^Uy_{ij})$, $(h_W)^*=(A^Wy_{ij},A^W(\{y_{lk},x_{p,q},z_{u,v}\}),B^Wy_{ij})$, and all other components of $(\operatorname{in}(\mathcal{L}))^*$ that involve y_{ij}. For $(h_X)^*$ that does not involve y_{ij} we leave as it is and similarly for others that do not involve y_{ij}.

Claim: There is a filtration from $\operatorname{in}(\mathcal{L})^*$ to an ideal

$$I_{y_{ij}}=(h_X)^*\cap(A^Y,B^Y)\cap(h_g)^*\cap(A^U,B^U)\cap...\cap(A^W,B^W)$$

such that the quotients are the form P/PL where P is an ideal and L is an ideal generated by variables such that they are a regular sequence modulo P.

With this claim, we can continue picking another variable and reduce to a bigger ideal that does not involve the new variable. We can continue the process until we reach an ideal that does not involve any $z_{i,n}$, or $x_{i,n}$ or $y_{i,n}$. This ideal is I_{n-1}.

Proof of claim: Without lost of generality, we just need to show: there is a filtration from $(A^Y(y_{ij},\{a_l^Y\}),B^Yy_{ij})\cap(A^U(y_{ij},\{a_l^U\}),B^Uy_{ij}))$ to $(A^Y,B^Y)\cap(A^U,B^U)$ with quotients are the form P/PL as above. For convenience, we write $A^Y(\{a_l^Y\})=A^YC$ and $A^U(\{a_l^U\})=A^UD$. Then we look at the following filtration:

$$\begin{aligned}
&(A^Yy_{ij},A^YC,B^Yy_{ij})\cap(A^Uy_{ij},A^UD,B^Uy_{ij})\\
=\;&((A^Y\cap A^U)y_{ij},(A^Y\cap A^U)Cy_{ij},(B^Y\cap A^U)y_{ij},(A^Y\cap A^U)CD,\\
&(A^Y\cap B^U)y_{ij},(B^Y\cap B^U)y_{ij}) \qquad =: J_0
\end{aligned}$$

$$\begin{array}{rcl}
\subset & ((A^Y \cap A^U)y_{ij}, (A^Y \cap A^U)C, (B^Y \cap A^U)y_{ij}, (A^Y \cap B^U)y_{ij}, (B^Y \cap B^U)y_{ij}) & =: J_1 \\
\subset & ((A^Y \cap A^U), (B^Y \cap A^U)y_{ij}, (A^Y \cap B^U)y_{ij}, (B^Y \cap B^U)y_{ij}) & =: J_2 \\
\subset & ((A^Y \cap A^U), (B^Y \cap A^U), (A^Y \cap B^U)y_{ij}, (B^Y \cap B^U)y_{ij}) & =: J_3 \\
\subset & ((A^Y \cap A^U), (B^Y \cap A^U), (A^Y \cap B^U), (B^Y \cap B^U)y_{ij}) & =: J_4 \\
\subset & ((A^Y \cap A^U), (B^Y \cap A^U), (A^Y \cap B^U), (B^Y \cap B^U)) = (A^Y, B^Y) \cap (A^U, B^U) & =: J_5.
\end{array}$$

We have

$$\begin{array}{rc}
 & J_1/J_0 \\
= & (A^Y \cap A^U)C/((A^Y \cap A^U)CD, (A^Y \cap A^U)C \cap (A^Y \cap A^U)y_{ij}, (A^Y \cap A^U)C \cap \\
 & (B^Y \cap A^U)y_{ij}, (A^Y \cap A^U)C \cap (A^Y \cap B^U)y_{ij}, (A^Y \cap A^U)C \cap (B^Y \cap B^U)y_{ij}) \\
= & (A^Y \cap A^U)C/((A^Y \cap A^U)C(D, y_{ij}), (A^Y \cap B^Y) \cap A^U C y_{ij}, (A^Y \cap A^U \cap B^U)C y_{ij}, \\
 & (A^Y \cap B^Y) \cap (A^U \cap B^U)C y_{ij}) \\
= & (A^Y \cap A^U)C/((A^Y \cap A^U)C(D, y_{ij}), (A^Y a^Y) \cap A^U C y_{ij}, (A^Y \cap A^U a^U)C y_{ij}, \\
 & (A^Y a^Y) \cap (A^U a^U)C y_{ij}) \\
= & (A^Y \cap A^U)C/((A^Y \cap A^U)C(D, y_{ij})).
\end{array}$$

Hence $\operatorname{reg}(J_1/J_0) = \operatorname{reg}(A^Y \cap A^U)C$. Also

$$\begin{array}{rc}
 & J_2/J_1 \\
= & (A^Y \cap A^U)/((A^Y \cap A^U)(C, y_{ij}), (A^Y \cap A^U) \cap (B^Y \cap A^U)y_{ij}, (A^Y \cap A^U) \cap \\
 & (A^Y \cap B^U)y_{ij}, (A^Y \cap A^U) \cap (B^Y \cap B^U)y_{ij}) \\
= & (A^Y \cap A^U)/((A^Y \cap A^U)(C, y_{ij}), (A^Y \cap B^Y \cap A^U)y_{ij}, (A^Y \cap A^U \cap B^U)y_{ij}, \\
 & (A^Y \cap B^Y) \cap (A^U \cap B^U)y_{ij})
\end{array}$$

$$\begin{array}{rc}
= & (A^Y \cap A^U)/((A^Y \cap A^U)(C, y_{ij}), (A^Y a^Y \cap A^U)y_{ij}, (A^Y \cap A^U a^U)y_{ij}, \\
 & (A^Y a^Y) \cap (A^U a^U)y_{ij}) \\
= & (A^Y \cap A^U)/((A^Y \cap A^U)(C, y_{ij})).
\end{array}$$

Hence $\operatorname{reg}(J_2/J_1) = \operatorname{reg}(A^Y \cap A^U)$. Similarly for

$$\begin{array}{rc}
 & J_3/J_2 \\
= & (B^Y \cap A^U)/((B^Y \cap A^U)y_{ij}, (B^Y \cap A^U) \cap (A^Y \cap A^U), (B^Y \cap A^U) \cap (A^Y \cap B^U)y_{ij}, \\
 & (B^Y \cap A^U) \cap (B^Y \cap B^U)y_{ij}) \\
= & (B^Y \cap A^U)/((B^Y \cap A^U)y_{ij}, (B^Y \cap A^Y \cap A^U), (B^Y \cap A^Y) \cap (A^U \cap B^U)y_{ij}, \\
 & (B^Y \cap A^U \cap B^U)y_{ij}) \\
= & (B^Y \cap A^U)/((B^Y \cap A^U)y_{ij}, (B^Y(\{b_i^Y\}) \cap A^U), (B^Y(\{b_i^Y\})) \cap (A^U a^U)y_{ij}, \\
 & (B^Y \cap A^U a^U)y_{ij}) \\
= & (B^Y \cap A^U)/((B^Y \cap A^U)(\{b_i^Y\}, y_{ij})).
\end{array}$$

Thus $\operatorname{reg}(J_3/J_2) = \operatorname{reg}(B^Y \cap A^U)$. Similarly for

$$\begin{aligned} & J_4/J_3 \\ = \;& (A^Y \cap B^U)/((A^Y \cap B^U)y_{ij}, (A^Y \cap B^U) \cap (A^Y \cap A^U), (A^Y \cap B^U) \cap (B^Y \cap A^U), \\ & (A^Y \cap B^U) \cap (B^Y \cap B^U)y_{ij}) \\ = \;& (A^Y \cap B^U)/((A^Y \cap B^U)y_{ij}, (A^Y \cap B^U(\{b_i^Y\})), (A^Y a^Y \cap B^U(\{b_i^Y\})), \\ & (A^Y a^Y \cap B^U y_{ij})) \\ = \;& (A^Y \cap B^U)/((A^Y \cap B^U)(y_{ij}, \{b_i^Y\})). \end{aligned}$$

Therefore $\operatorname{reg}(J_4/J_3) = \operatorname{reg}(A^Y \cap B^U)$. Finally,

$$\begin{aligned} & J_5/J_4 \\ = \;& (B^Y \cap B^U)/((B^Y \cap B^U)y_{ij}, (B^Y \cap B^U) \cap (A^Y \cap A^U), (B^Y \cap B^U) \cap \\ & (B^Y \cap A^U), (B^Y \cap B^U) \cap (A^Y \cap B^U)) \\ = \;& (B^Y \cap B^U)/((B^Y \cap B^U)y_{ij}, (B^Y(\{b_i^Y\}) \cap B^U(\{b_i^U\})), \\ & (B^Y \cap B^U(\{b_i^U\})), (B^Y(\{b_i^Y\}) \cap B^U)) \\ = \;& (B^Y \cap B^U)/((B^Y \cap B^U)(y_{ij}, \{b_i^Y\}, \{b_i^U\}). \end{aligned}$$

We obtain $\operatorname{reg}(J_5/J_4) = \operatorname{reg}(B^Y \cap B^U)$.

Since by induction hypothesis we have

$$\operatorname{reg}(A^Y, B^Y) \cap (A^U, B^U) = d = \operatorname{d}(A^Y, B^Y) \cap (A^U, B^U),$$

it follows that

$$\operatorname{d}(A^Y \cap A^U) = \operatorname{d}(B^Y \cap A^U) = \operatorname{d}(A^Y \cap B^U) = \operatorname{d}(B^Y \cap B^U) = d$$

and

$$\operatorname{reg}(A^Y \cap A^U) = \operatorname{reg}(B^Y \cap A^U) = \operatorname{reg}(A^Y \cap B^U) = \operatorname{reg}(B^Y \cap B^U) = d.$$

Also notice that $\operatorname{reg}(A^Y \cap A^U)C \geq \operatorname{d}(A^Y \cap A^U)C \geq d+1$. We use the regularity of the quotients of the filtration. to obtain the regularity of J_0. We have $\operatorname{reg}J_4 = \operatorname{reg}(J_5) + 1 = d+1$ and $\operatorname{reg}J_3 = d+1 = \operatorname{reg}J_2 = d+1 = \operatorname{reg}J_1 = d+1$. Notice $\operatorname{reg}J_0 \geq \deg J_0 \geq \deg J_1 + 1 = d+1$. We will show $\operatorname{reg}J_0 \leq d+1$, hence $\operatorname{reg}J_0 = \deg J_0 = d+1$. Assume $\operatorname{reg}J_0 > d+2$, then $\operatorname{reg}J_1 = \max\{\operatorname{reg}J_0, \operatorname{reg}J_1/J_0 = \operatorname{reg}(A^Y \cap A^U)C\} > d+2$, a contradiction. Hence $\operatorname{reg}J_0 = d+1 = \deg J_0$. This completes the proof of the claim.

Since $s_1 \geq s_2$, we assume $s_1 = m$ and we observe that $(\operatorname{in}(\mathcal{L}))^*$ does not involve y_{in} for $i > 2$ and z_{mn}. If $t_2 < n$, then $(\operatorname{in}(\mathcal{L}))^*$ does not involve y_{in} for all i. By using the claim above, we can find a filtration starting from $(\operatorname{in}(\mathcal{L}))^*$ to an ideal $J_{y_{1n}}$, where y_{1n} is not a factor of the minimal monomial generators of $J_{y_{1n}}$. Then we continue the filtration to an ideal J_{2n} such that y_{2n} is not a factor of the minimal monomial generators of $J_{y_{2n}}$. We need those two steps when $t_2 = n$, otherwise we skip those steps. The next step is to look at $z_{m-1,n}$ and find a filtration until an ideal $J_{z,_{m-1},n}$ such that $z_{m-1,n}$ is not a factor of $J_{z_{m-1,n}}$. Next we consider $z_{m-2,n}$ and continue to z_{1n}. Finally, we consider x_{1n}. Then we will get the ideal I_{n-1}. We will have a filtration as follow:

$$I_n \subset J_{y_{1n}} \subset J_{y_{2n}} \subset J_{m-1,n} \subset J_{m-2,n} \subset .... \subset J_{x_{1n}} = I_{n-1}.$$

Hence by using a similar argument as in the proof of the claim, we have

$$\begin{aligned}
\mathrm{reg} I_n &= \mathrm{reg} I_{n-1} + 2 + m - 1 + 1 \\
&= m(n-1) - 1 + t_1 - 1 - (s_1 - 1) + t_2 - 1 - (s_2 - 1) + m + 2 \\
&= mn - 1 + t_1 - (s_1 - 1) + t_2 - (s_2 - 1) \\
&= \mathrm{d} I_{n-1} + 2 + m - 1 + 1 \\
&= \mathrm{d} I_n.
\end{aligned}$$

This complete the proof of this Lemma.

□

We are now ready to prove Theorem 1.

Proof of Theorem 1: We know that $k[X,Y,Z]/(\mathrm{in}(\mathcal{K})$ is Cohen-Macaulay by Lemma 21 and Theorem 3. Hence $\mathcal{R}(\mathbb{D}) = k[X,Y,Z]/\mathcal{K}$ is Cohen-Macaulay [E]. □

References

[B-H] W. Bruns and J. Herzog, Cohen-Macaulay rings, Cambridge Studies in Advanced Mathematics 39, Cambridge University Press, Cambridge, 1993.

[E-R] J. Eagon and V. Reiner, Resolutions of Stanley-Reisner rings and Alexander duality, J. Pure Appl. Algebra 130 (1998), 165-175.

[E] Commutative algebra with a view toward algebraic geometry, Graduate Texts in Mathematics 150, Springer-Verlag, New York, 1995.

[G-S] D. Grayson and M. Stillman, Macaulay 2, A computer algebra system for computing in algebraic geometry and commutative algebra, available through anonymous ftp from http://www.math.uiuc.edu/Macaulay2.

[G-P-S] G. Greuel, G. Pfister, and H. Schonemann, Singular 3.1.0, A computer algebra system for polynomial computations, available through http://www.singular.uni-kl.de.

[H-T-Z] J. Herzog, Z. Tang and S. Zarzuela, Symmetric and Rees algebras of Koszul cycles and their Groebner bases, Manuscripta Math. 112 (2003), 489-509.

[L] K-N Lin, Rees Algebras of Diagonal Ideals, preprint.

[M] A. Micali, Sur les algebres universelles, Ann. Inst. Fourier (Grenoble) 14 (1964), fasc. 2, 33–87.

[S-S-U] A. Simis, K. Smith and B. Ulrich, An algebraic proof of Zak's inequality for the dimension of the Gauss image, Math. Z. 241 (2002), 871-881.

[S-U] A. Simis and B. Ulrich, On the ideal of an embedded join, J. Algebra 226 (2000), 1-14.

[S-S] B. Sturmfels and S. Sullivant, Combinatorial secant varieties, Pure Appl. Math. 2 (2006), 867-891.

Department of Mathematics,
University of California, Riverside, CA 92521, USA
e-mail: linkuei@ucr.edu